

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
4 March 2004 (04.03.2004)

PCT

(10) International Publication Number
WO 2004/018451 A1

(51) International Patent Classification⁷: **C07D 401/04**,
401/14, A61K 31/50

(74) Agent: **WILD, Robert**; Altana Pharma AG, Byk-Gulden-
Strasse 2, 78467 Konstanz (DE).

(21) International Application Number:
PCT/EP2003/008677

(81) Designated States (*national*): AE, AL, AU, BA, BR, CA,
CN, CO, DZ, EC, GE, HR, ID, IL, IN, IS, JP, KR, LT, LV,
MA, MK, MX, NO, NZ, PH, PL, SG, TN, UA, US, VN,
YU, ZA, ZW.

(22) International Filing Date: 6 August 2003 (06.08.2003)

(25) Filing Language: English

(84) Designated States (*regional*): Eurasian patent (AM, AZ,
BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE,
BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU,
IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR).

(26) Publication Language: English

(30) Priority Data:
02017976.8 10 August 2002 (10.08.2002) EP

Declarations under Rule 4.17:

(71) Applicant (*for all designated States except US*): **ALTANA
PHARMA AG** [DE/DE]; Byk-Gulden-Str. 2, 78467 Kon-
stanz (DE).

— as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AL, AU, BA, BR, CA, CN, CO, DZ, EC, GE, HR, ID, IL, IN,
IS, JP, KR, LT, LV, MA, MK, MX, NO, NZ, PH, PL, SG, TN,
UA, VN, YU, ZA, ZW, Eurasian patent (AM, AZ, BY, KG,
KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH,
CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU,
MC, NL, PT, RO, SE, SI, SK, TR)

(72) Inventors: **HATZELMANN, Armin**; Alter Wall 3, 78467
Konstanz (DE). **BARSIG, Johannes**; Bleichenweg 11,
78467 Konstanz (DE). **MARX, Degenhard**; Obere Reute
15, 78345 Moos (DE). **KLEY, Hans-Peter**; Im Weinberg
3b, 78476 Allensbach (DE). **CHRISTIAANS, Johannes**,
A. M.; Zevenwouden 233, NL-3524 CR Utrecht (NL).
MENGE, Wiro, M., P., B.; Pontanuslaan 11, NL-6821
HM Arnhem (NL).

— as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for all designations
of inventorship (Rule 4.17(iv)) for US only

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): **STERK, Geert, Jan**
[NL/NL]; Stadhouderslaan 38, NL-3583 JJ Utrecht (NL).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **PYRIDAZINONE-DERIVATIVES AS PDE4 INHIBITORS**

(57) Abstract: The compounds of a certain formula (1), in which the given substituents have the meanings as given in the description, are novel effective PDE4 or PDE3/4 inhibitors.

WO 2004/018451 A1

PYRIDAZINONE-DERIVATIVES AS PDE4 INHIBITORS

Field of application of the invention

The invention relates to novel pyridazinone-derivatives, which are used in the pharmaceutical industry for the production of pharmaceutical compositions.

Known technical background

International Patent Applications WO98/31674 (= USP 6,103,718), WO99/31071, WO99/31090, WO99/47505 (= USP 6,255,303), WO01/19818, WO01/30766, WO01/30777, WO01/94319, WO02/064584, WO02/085885 and WO02/085906 disclose phthalazinone derivatives having PDE4 inhibitory properties. In the International Patent Application WO94/12461 and in the European Patent Application EP 0 763 534 3-aryl-pyridazin-6-one and arylalkyl-diazinone derivatives are described as selective PDE4 inhibitors. International Patent Application WO93/07146 (= USP 5,716,954) discloses benzo and pyrido pyridazinone and pyridazinthione compounds with PDE4 inhibiting activity.


In the Journal of Medicinal Chemistry, Vol. 33, No. 6, 1990, pp. 1735-1741 1,4-Bis(3-oxo-2,3-dihydropyridazin-6-yl)benzene derivatives are described as potent phosphodiesterase inhibitors and inodilators. In the Journal of Medicinal Chemistry Vol. 45 No.12, 2002, pp. 2520-2525, 2526-2533 and in Vol. 44, No. 16, 2001, pp. 2511-2522 and pp. 2523-2535 phthalazinone derivatives are described as selective PDE4 inhibitors.

Description of the invention

It has now been found that the pyridazinone-derivatives, which are described in greater details below, have surprising and particularly advantageous properties.

The invention thus relates to compounds of formula 1

- 2 -





in which

R1 is hydrogen or 1-4C-alkyl,

R2 is hydrogen or 1-4C-alkyl,

R3 represents a phenyl derivative of formulae (a) or (b)



wherein

R4 is 1-4C-alkoxy or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R5 is 1-8C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy, or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R6 is 1-4C-alkoxy, 3-5C-cycloalkoxy, 3-5C-cycloalkylmethoxy, or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R7 is 1-4C-alkyl and

R8 is hydrogen or 1-4C-alkyl,

or wherein

R7 and R8 together and with inclusion of the two carbon atoms, to which they are bonded, form a spiro-linked 5-, 6- or 7-membered hydrocarbon ring, optionally interrupted by an oxygen or sulphur atom,

R9 is 1-4C-alkyl, $-S(O)_2-R_{10}$, $-S(O)_2-(CH_2)_n-R_{11}$, $-(CH_2)_m-S(O)_2-R_{12}$, $-C(O)R_{13}$, $-C(O)-(CH_2)_n-R_{14}$, $-(CH_2)_m-C(O)-R_{15}$, Aryl1 or (Aryl2)-1-4C-alkyl,

R10 is 1-4C-alkyl, 5-dimethylaminonaphthalin-1-yl, $-N(R_{16})R_{17}$, thiophenyl, phenyl or phenyl substituted by R18 and/or R19,

R11 is phenyl or $-N(R_{16})R_{17}$,

R12 is $-N(R_{16})R_{17}$,

R13 is 1-4C-alkyl, hydroxycarbonyl-1-4C-alkyl, phenyl, 2,4,6-trichlorophenyl, pyridyl, 4-ethyl-piperazin-2,3-dion-1-yl or $-N(R_{16})R_{17}$,

R14 is -N(R16)R17,


R15 is -N(R16)R17, phenyl or phenyl substituted by R18 and/or R19 and/or R20,

R16 is hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkylmethyl, phenyl or phenyl substituted by R18 and/or R19 and/or R20,

R17 1-7C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkylmethyl, phenyl or phenyl substituted by R18 and/or R19 and/or R20,

or

R16 and R17 together and with inclusion of the nitrogen atom to which they are bonded, form a 4-morpholinyl-, 1-pyrrolidinyl-, 1-piperidinyl-, 1-hexahydroazepino- or a 1-piperazinyl-ring of formula (c)



wherein

R21 is 1-4C-alkyl, pyrid-4-yl, pyrid-4-ylmethyl, 2-methoxyphenyl, 1,1-diphenylmethyl, dimethyl-amino-1-4C-alkyl, dimethylaminocarbonylmethyl, N-methyl-piperidin-4-yl, 4-morpholino-ethyl or tetrahydrofuran-2-ylmethyl,

R18 is halogen, nitro, cyano, carboxyl, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino, mono- or di-1-4C-alkylamino, aminocarbonyl, 1-4C-alkylcarbonylamino or mono- or di-1-4C-alkylaminocarbonyl,

R19 is halogen, amino, nitro, 1-4C-alkyl or 1-4C-alkoxy,

R20 is halogen,

Aryl1 is pyrimidin-2-yl, thieno-[2,3-d]pyrimidin-4-yl, 1-methyl-1H-pyrazolo-[3,4-d]pyrimidin-4-yl, thiazolyl, imidazolyl, furanyl, pyridyl, phenyl or phenyl substituted by R18 and/or R19,

Aryl2 is pyridyl, phenyl, phenyl substituted by R18 and/or R19, 2-oxo-2H-chromen-7-yl or 4-(1,2,3-thiadiazol-4-yl)phenyl,

n is an integer from 1 to 4,

m is an integer from 1 to 4,

and the salts of these compounds.

1-4C-Alkyl is a straight-chain or branched alkyl radical having 1 to 4 carbon atoms. Examples are the butyl, isobutyl, sec-butyl, tert-butyl, propyl, isopropyl, ethyl and methyl radicals.

1-7C-Alkyl is a straight-chain or branched alkyl radical having 1 to 7 carbon atoms. Examples are the heptyl, isoheptyl (5-methylhexyl), hexyl, isohexyl (4-methylpentyl), neoheptyl (3,3-dimethylbutyl), pentyl, isopentyl (3-methylbutyl), neopentyl (2,2-dimethylpropyl), butyl, isobutyl, sec-butyl, tert-butyl, propyl, isopropyl, ethyl and methyl radicals.

1-4C-Alkoxy is a radical which, in addition to the oxygen atom, contains a straight-chain or branched alkyl radical having 1 to 4 carbon atoms. Alkoxy radicals having 1 to 4 carbon atoms which may be mentioned in this context are, for example, the butoxy, isobutoxy, sec-butoxy, tert-butoxy, propoxy, isopropoxy, ethoxy and methoxy radicals.

1-8C-Alkoxy is a radical which, in addition to the oxygen atom, contains a straight-chain or branched alkyl radical having 1 to 8 carbon atoms. Alkoxy radicals having 1 to 8 carbon atoms which may be mentioned in this context are, for example, the octyloxy, heptyloxy, isoheptyloxy (5-methylhexyloxy), hexyloxy, isohexyloxy (4-methylpentyloxy), neoheptyloxy (3,3-dimethylbutoxy), pentyloxy, isopentyloxy (3-methylbutoxy), neopentyloxy (2,2-dimethylpropoxy), butoxy, isobutoxy, sec-butoxy, tert-butoxy, propoxy, isopropoxy, ethoxy and methoxy radicals.

Halogen within the meaning of the present invention is bromine, chlorine or fluorine.

3-7C-Cycloalkoxy stands for cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy or cycloheptyloxy, of which cyclopropyloxy, cyclobutyloxy and cyclopentyloxy are preferred.

3-7C-Cycloalkylmethoxy stands for cyclopropylmethoxy, cyclobutylmethoxy, cyclopentylmethoxy, cyclohexylmethoxy or cycloheptylmethoxy, of which cyclopropylmethoxy, cyclobutylmethoxy and cyclopentylmethoxy are preferred.

3-5C-Cycloalkoxy stands for cyclopropyloxy, cyclobutyloxy and cyclopentyloxy.

3-5C-Cycloalkylmethoxy stands for cyclopropylmethoxy, cyclobutylmethoxy and cyclopentylmethoxy.

3-7C-Cycloalkyl stands for cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl, of which cyclopropyl and cyclopentyl are preferred

3-7C-Cycloalkylmethyl stands for cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl or cycloheptylmethyl.

1-4C-Alkoxy which is completely or predominantly substituted by fluorine is, for example, the 2,2,3,3,3-pentafluoropropoxy, the perfluoroethoxy, the 1,2,2-trifluoroethoxy and in particular the 1,1,2,2-tetrafluoroethoxy, the 2,2,2-trifluoroethoxy, the trifluoromethoxy and the difluoromethoxy radical, of which the difluoromethoxy radical is preferred. "Predominantly" in this connection means that more than half of the hydrogen atoms of the 1-4C-alkoxy group are replaced by fluorine atoms.

As spiro-linked 5-, 6- or 7-membered hydrocarbon rings, optionally interrupted by an oxygen or sulphur atom, may be mentioned the cyclopentane, cyclohexane, cycloheptane, tetrahydrofuran, tetrahydropyran and the tetrahydrothiophen ring.

1-4C-Alkoxy carbonyl is a carbonyl group to which one of the abovementioned 1-4C-alkoxy radicals is bonded. Examples are the methoxycarbonyl [$\text{CH}_3\text{O}-\text{C}(\text{O})-$] and the ethoxycarbonyl [$\text{CH}_3\text{CH}_2\text{O}-\text{C}(\text{O})-$] radical.

An 1-4C-Alkylcarbonylamino radical is, for example, the propionylamino [$\text{C}_3\text{H}_7\text{C}(\text{O})\text{NH}-$] and the acetyl-amino radical [$\text{CH}_3\text{C}(\text{O})\text{NH}-$].

Mono- or Di-1-4C-alkylamino radicals contain in addition to the nitrogen atom, one or two of the abovementioned 1-4C-alkyl radicals. Preferred are the di-1-4C-alkylamino radicals, especially the dimethylamino, the diethylamino and the diisopropylamino radical.

Mono- or Di-1-4C-alkylaminocarbonyl radicals contain in addition to the carbonyl group one of the abovementioned mono- or di-1-4C-alkylamino radicals. Examples which may be mentioned are the N-methyl-, the N,N-dimethyl-, the N-ethyl-, the N-propyl-, the N,N-diethyl- and the N-isopropylaminocarbonyl radical.

(Aryl₂)-1-4C-alkyl radicals stand for one of the abovementioned 1-4C-alkyl radicals substituted by an Aryl₂ radical. Examples which may be mentioned are the pyrid-3-ylmethyl, pyrid-4-ylmethyl or benzyl radical.

Hydroxycarbonyl-1-4C-alkyl stand for one of the abovementioned 1-4C-alkyl radicals substituted by a hydroxycarbonyl (carboxyl) radical.

Dimethylamino-1-4C-alkyl radicals stand for one of the abovementioned 1-4C-alkyl radicals substituted by a dimethylamino radical.

Suitable salts for compounds of the formula 1 are all acid addition salts. Particular mention may be made of the pharmacologically tolerable inorganic and organic acids customarily used in pharmacy. Those suitable are water-soluble and water-insoluble acid addition salts with acids such as, for example, hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid, sulphuric acid, acetic acid, citric acid, D-gluconic acid, benzoic acid, 2-(4-hydroxybenzoyl)benzoic acid, butyric acid, sulphosalicylic acid, maleic acid, lauric acid, malic acid, fumaric acid, succinic acid, oxalic acid, tartaric acid, embonic acid, stearic acid, toluenesulphonic acid, methanesulphonic acid or 3-hydroxy-2-naphthoic acid, the acids being employed in salt preparation - depending on whether a mono- or polybasic acid is concerned and depending on which salt is desired - in an equimolar quantitative ratio or one differing therefrom.

Pharmacologically intolerable salts, which can be obtained, for example, as process products during the preparation of the compounds according to the invention on an industrial scale, are converted into pharmacologically tolerable salts by processes known to the person skilled in the art.


According to expert's knowledge the compounds of the invention as well as their salts may contain, e.g. when isolated in crystalline form, varying amounts of solvents. Included within the scope of the invention are therefore all solvates and in particular all hydrates of the compounds of formula 1 as well as all solvates and in particular all hydrates of the salts of the compounds of formula 1.

Compounds of formula 1 to be emphasized are those in which

R1 is hydrogen,

R2 is hydrogen or 1-4C-alkyl,

R3 represents a phenyl derivative of formulae (a) or (b)



wherein

R4 is 1-2C-alkoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,

R5 is 1-4C-alkoxy,

R6 is 1-2C-alkoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,

R7 is methyl and

R8 is hydrogen,

or wherein

R7 and R8 together and with inclusion of the two carbon atoms, to which they are bonded, form a spiro-linked cyclopentane, cyclohexane, tetrahydrofurane or tetrahydropyran ring,

R9 is 1-4C-alkyl, -S(O)₂-R10, -S(O)₂-(CH₂)_n-R11, -C(O)R13, -C(O)-(CH₂)_n-R14, -(CH₂)_m-C(O)-R15 or (Aryl₂)-1-4C-alkyl,

R10 is 1-4C-alkyl, 5-dimethylaminonaphthalin-1-yl, -N(R16)R17, thiophenyl, phenyl or phenyl substituted by R18 and/or R19,

R11 is phenyl,

R13 is 1-4C-alkyl, phenyl, pyridyl, 2,4,6-trichlorophenyl, 4-ethyl-piperazin-2,3-dion-1-yl or -N(R16)R17,

R14 is -N(R16)R17,


R15 is -N(R16)R17, phenyl or phenyl substituted by R18 and/or R19,

R16 is hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl or 3-7C-cycloalkylmethyl,

R17 is 1-7C-alkyl, 3-7C-cycloalkyl or 3-7C-cycloalkylmethyl,

or

R16 and R17 together and with inclusion of the nitrogen atom to which they are bonded, form a 4-morpholinyl-, 1-pyrrolidinyl-, 1-piperidinyl-, 1-hexahydroazepino- or a 1-piperazinyl-ring of formula (c)



wherein

R21 is 1-4C-alkyl, pyrid-4-yl, 2-methoxyphenyl, 1,1-diphenylmethyl or N-methyl-piperidin-4-yl,

R18 is halogen, nitro, cyano, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine or 1-4C-alkoxycarbonyl,

R19 is halogen, amino, nitro, 1-4C-alkyl or 1-4C-alkoxy,

Aryl2 is pyridyl, phenyl or phenyl substituted by R18 and/or R19,

m is an integer from 1 to 2,

n is an integer from 1 to 2,


and the salts of these compounds.

Preferred compounds of formula 1 are those, in which

R1 is hydrogen,

R2 is hydrogen or methyl,

R3 represents a phenyl derivative of formula (a)



wherein

R4 is 1-2C-alkoxy,

R5 is 1-2C-alkoxy,

R9 is -S(O)₂-R10, -S(O)₂-(CH₂)_n-R11, -C(O)R13, -C(O)-(CH₂)_n-R14, -(CH₂)_m-C(O)-R15 or (Aryl2)-1-2C-alkyl,

R10 is 1-4C-alkyl, 5-dimethylaminonaphthalin-1-yl, -N(R16)R17, thiophenyl, phenyl or phenyl substituted by R18 and/or R19,

R11 is phenyl,

R13 is 1-4C-alkyl, phenyl, 2,4,6-trichlorophenyl, pyridyl, 4-ethyl-piperazin-2,3-dion-1-yl or -N(R16)R17,

R14 is -N(R16)R17,


R15 is -N(R16)R17,

R16 is hydrogen or 1-4C-alkyl,

R17 is 1-4C-alkyl,

or

R16 and R17 together and with inclusion of the nitrogen atom to which they are bonded, form a 4-morpholinyl-, 1-pyrrolidinyl-, 1-piperidinyl- or a 1-piperazinyl-ring of formula (c)



wherein

R21 is 1-4C-alkyl, pyrid-4-yl, 2-methoxyphenyl or 1,1-diphenylmethyl,

R18 is halogen, cyano, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R19 is 1-4C-alkyl or 1-4C-alkoxy,

Aryl2 is pyridyl or phenyl,

m is 1,

n is 1,


and the salts of these compounds.

Particularly preferred compounds of formula 1 are those in which

R1 is hydrogen,

R2 is hydrogen or methyl,

R3 represents a phenyl derivative of formula (a)



wherein

R4 is methoxy,

R5 is methoxy and

R9 is acetyl, morpholin-4-ylcarbonyl, pyridin-3-ylmethyl, 4-ethyl-piperazin-2,3-dion-1-ylcarbonyl, 4-methylpiperazin-1-ylcarbonyl, 5-dimethylamino-naphthalene-1-sulfonyl, 2-(morpholin-4-yl)-2-oxo-ethyl, 4-methylbenzenesulfonyl, methylsulfonyl, 4-chlorobenzenesulfonyl, benzy sulfonyl, 4-methoxybenzenesulfonyl, benzenesulfonyl, 2,5-dimethoxybenzenesulfonyl, 2-cyanobenzenesulfonyl, thiophen-2-ylsulfonyl, 2-fluorobenzenesulfonyl, 2-trifluoromethoxy-


benzenesulfonyl, dimethylaminosulfonyl, benzoyl, pyridin-3-ylcarbonyl, 2,4,6-trichlorobenzenecarbonyl, tert-butylaminocarbonyl, dimethylaminocarbonylmethyl, 2-(4-methyl-piperazin-1-yl)-2-oxo-ethyl, 2-(4-pyridin-4-ylpiperazin-1-yl)ethanoyl, 2-[4-(2-methoxyphenyl)piperazin-1-yl]ethanoyl or 2-[4-(1,1-diphenylmethyl)piperazin-1-yl]ethanoyl, and the salts of these compounds.

An embodiment (embodiment A) of the compounds of formula 1 are those in which

R1 is hydrogen or 1-4C-alkyl,

R2 is hydrogen or 1-4C-alkyl,

R3 represents a phenyl derivative of formulae (a) or (b)



wherein

R4 is 1-4C-alkoxy or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R5 is 1-8C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy, or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R6 is 1-4C-alkoxy, 3-5C-cycloalkoxy, 3-5C-cycloalkylmethoxy, or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R7 is 1-4C-alkyl and

R8 is hydrogen or 1-4C-alkyl,

or wherein

R7 and R8 together and with inclusion of the two carbon atoms, to which they are bonded, form a spiro-linked 5-, 6- or 7-membered hydrocarbon ring, optionally interrupted by an oxygen or sulphur atom,

R9 is $-S(O)_2-R_{10}$, $-S(O)_2-(CH_2)_n-R_{11}$ or $-(CH_2)_m-S(O)_2-R_{12}$,

R10 is 1-4C-alkyl, 5-dimethylaminonaphthalin-1-yl, $-N(R_{16})R_{17}$, thiophenyl, phenyl or phenyl substituted by R18 and/or R19,

R11 is phenyl or $-N(R_{16})R_{17}$,


R12 is $-N(R_{16})R_{17}$,

R16 is hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkylmethyl, phenyl or phenyl substituted by R18 and/or R19 and/or R20,

R17 1-7C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkylmethyl, phenyl or phenyl substituted by R18 and/or R19 and/or R20,

or

R16 and R17 together and with inclusion of the nitrogen atom to which they are bonded, form a 4-morpholinyl-, 1-pyrrolidinyl-, 1-piperidinyl-, 1-hexahydroazepino- or a 1-piperazinyl-ring of formula (c)



wherein

R21 is 1-4C-alkyl, pyrid-4-yl, pyrid-4-ylmethyl, 2-methoxyphenyl, 1,1-diphenylmethyl, dimethyl-amino-1-4C-alkyl, dimethylaminocarbonylmethyl, N-methyl-piperidin-4-yl, 4-morpholino-ethyl or tetrahydrofuran-2-ylmethyl,

R18 is halogen, nitro, cyano, carboxyl, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino, mono-or di-1-4C-alkylamino, aminocarbonyl, 1-4C-alkylcarbonylamino or mono-or di-1-4C-alkylaminocarbonyl,

R19 is halogen, amino, nitro, 1-4C-alkyl or 1-4C-alkoxy,

R20 is halogen,

n is an integer from 1 to 4,

m is an integer from 1 to 4,


and the salts of these compounds.

Preferred compounds of formula 1 of embodiment A are those in which

R1 is hydrogen,

R2 is hydrogen or methyl,

R3 represents a phenyl derivative of formula (a)



wherein

R4 is 1-2C-alkoxy,

R5 is 1-2C-alkoxy,

R9 is -S(O)₂-R10 or -S(O)₂-(CH₂)_n-R11,

R10 is 1-4C-alkyl, 5-dimethylaminonaphthalin-1-yl, -N(R16)R17, thiophenyl, phenyl or phenyl substituted by R18 and/or R19,

R11 is phenyl,

R16 is hydrogen or 1-4C-alkyl,

R17 is 1-4C-alkyl,

R18 is halogen, cyano, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R19 is 1-4C-alkyl or 1-4C-alkoxy,

n is 1,


and the salts of these compounds.

Particularly preferred compounds of formula 1 of embodiment A are those in which

R1 is hydrogen,

R2 is hydrogen or methyl,

R3 represents a phenyl derivative of formula (a)



wherein

R4 is methoxy,

R5 is methoxy and

R9 is 5-dimethylamino-naphthalene-1-sulfonyl, 4-methylbenzenesulfonyl, methylsulfonyl, 4-chlorobenzenesulfonyl, benzylsulfonyl, 4-methoxybenzenesulfonyl, benzenesulfonyl, 2,5-dimethoxybenzenesulfonyl, 2-cyanobenzenesulfonyl, thiophen-2-ylsulfonyl, 2-fluorobenzenesulfonyl, 2-trifluoromethoxy-benzenesulfonyl or dimethylaminosulfonyl,



and the salts of these compounds.

Another embodiment (embodiment B) of the compounds of formula 1 are those in which

R1 is hydrogen or 1-4C-alkyl,

R2 is hydrogen or 1-4C-alkyl,

R3 represents a phenyl derivative of formulae (a) or (b)



wherein

R4 is 1-4C-alkoxy or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R5 is 1-8C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy, or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R6 is 1-4C-alkoxy, 3-5C-cycloalkoxy, 3-5C-cycloalkylmethoxy, or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R7 is 1-4C-alkyl and

R8 is hydrogen or 1-4C-alkyl,

or wherein

R7 and R8 together and with inclusion of the two carbon atoms, to which they are bonded, form a spiro-linked 5-, 6- or 7-membered hydrocarbon ring, optionally interrupted by an oxygen or sulphur atom,

R9 is -C(O)R13, -C(O)-(CH₂)_n-R14 or -(CH₂)_m-C(O)-R15,

R13 is 1-4C-alkyl, hydroxycarbonyl-1-4C-alkyl, phenyl, 2,4,6-trichlorophenyl, pyridyl, 4-ethyl-piperazin-2,3-dione-1-yl or -N(R16)R17,

R14 is -N(R16)R17,


R15 is -N(R16)R17, phenyl or phenyl substituted by R18 and/or R19 and/or R20,

R16 is hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkylmethyl, phenyl or phenyl substituted by R18 and/or R19 and/or R20,

R17 is 1-7C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkylmethyl, phenyl or phenyl substituted by R18 and/or R19 and/or R20,

or

R16 and R17 together and with inclusion of the nitrogen atom to which they are bonded, form a 4-morpholinyl-, 1-pyrrolidinyl-, 1-piperidinyl-, 1-hexahydroazepino- or a 1-piperazinyl-ring of formula (c)



wherein

R21 is 1-4C-alkyl, pyrid-4-yl, pyrid-4-ylmethyl, 2-methoxyphenyl, 1,1-diphenylmethyl, dimethyl-amino-1-4C-alkyl, dimethylaminocarbonylmethyl, N-methyl-piperidin-4-yl, 4-morpholino-ethyl or tetrahydrofuran-2-ylmethyl,

R18 is halogen, nitro, cyano, carboxyl, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino, mono- or di-1-4C-alkylamino, aminocarbonyl, 1-4C-alkylcarbonylamino or mono- or di-1-4C-alkylaminocarbonyl,

R19 is halogen, amino, nitro, 1-4C-alkyl or 1-4C-alkoxy,

R20 is halogen,

n is an integer from 1 to 4,

m is an integer from 1 to 4,


and the salts of these compounds.

Preferred compounds of formula 1 of embodiment B are those in which

R1 is hydrogen,

R2 is hydrogen or methyl,

R3 represents a phenyl derivative of formula (a)



wherein

R4 is 1-2C-alkoxy,

R5 is 1-2C-alkoxy,

R9 is -C(O)R13, -C(O)-(CH₂)_n-R14 or -(CH₂)_m-C(O)-R15,

R13 is 1-4C-alkyl, phenyl, 2,4,6-trichlorophenyl, pyridyl, 4-ethyl-piperazin-2,3-dion-1-yl or -N(R16)R17,

R14 is -N(R16)R17,


R15 is -N(R16)R17,

R16 is hydrogen or 1-4C-alkyl,

R17 is 1-4C-alkyl,

or

R16 and R17 together and with inclusion of the nitrogen atom to which they are bonded, form a 4-morpholinyl-, 1-pyrrolidinyl-, 1-piperidinyl- or a 1-piperazinyl-ring of formula (c)



wherein

R21 is 1-4C-alkyl, pyrid-4-yl, 2-methoxyphenyl or 1,1-diphenylmethyl,

m is 1,

n is 1,


and the salts of these compounds.

Particularly preferred compounds of formula 1 of embodiment B are those in which

R1 is hydrogen,

R2 is hydrogen or methyl,

R3 represents a phenyl derivative of formula (a)



wherein

R4 is methoxy,

R5 is methoxy and



R9 is acetyl, morpholin-4-ylcarbonyl, 4-ethyl-piperazin-2,3-dion-1-ylcarbonyl, 4-methylpiperazin-1-ylcarbonyl, 2-(morpholin-4-yl)-2-oxo-ethyl, benzoyl, pyridin-3-ylcarbonyl, 2,4,6-trichlorobenzenecarbonyl, tert-butylaminocarbonyl, dimethylaminocarbonylmethyl, 2-(4-methyl-piperazin-1-yl)-2-oxo-ethyl, 2-(4-pyridin-4-ylpiperazin-1-yl)ethanoyl, 2-[4-(2-methoxyphenyl)piperazin-1-yl]ethanoyl or 2-[4-(1,1-diphenylmethyl)piperazin-1-yl]ethanoyl, and the salts of these compounds.

Further compounds of formula 1 (embodiment C) are those in which

R1 is hydrogen or 1-4C-alkyl,

R2 is hydrogen or 1-4C-alkyl,

R3 represents a phenyl derivative of formulae (a) or (b)



wherein

R4 is 1-4C-alkoxy or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R5 is 1-8C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy, or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R6 is 1-4C-alkoxy, 3-5C-cycloalkoxy, 3-5C-cycloalkylmethoxy, or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,


R7 is 1-4C-alkyl and

R8 is hydrogen or 1-4C-alkyl,

or wherein

R7 and R8 together and with inclusion of the two carbon atoms, to which they are bonded, form a spiro-linked 5-, 6- or 7-membered hydrocarbon ring, optionally interrupted by an oxygen or sulphur atom,

- R9 is hydrogen, 1-4C-alkyl, $-S(O)_2-R_{10}$, $-S(O)_2-(CH_2)_n-R_{11}$, $-(CH_2)_m-S(O)_2-R_{12}$, $-C(O)R_{13}$, $-C(O)-(CH_2)_n-R_{14}$, $-(CH_2)_m-C(O)-R_{15}$, Aryl1 or (Aryl2)-1-4C-alkyl,
- R10 is 1-4C-alkyl, 5-dimethylaminonaphthalin-1-yl, $-N(R_{16})R_{17}$, phenyl or phenyl substituted by R18 and/or R19,
- R11 is $-N(R_{16})R_{17}$,
- R12 is $-N(R_{16})R_{17}$,
- R13 is 1-4C-alkyl, hydroxycarbonyl-1-4C-alkyl, phenyl, pyridyl, 4-ethyl-piperazin-2,3-dion-1-yl or $-N(R_{16})R_{17}$,
- R14 is $-N(R_{16})R_{17}$,
- R15 is $-N(R_{16})R_{17}$, phenyl or phenyl substituted by R18 and/or R19 and/or R20,
- R16 and R17 are independent from each other hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-methyl, phenyl or phenyl substituted by R18 and/or R19 and/or R20, or R16 and R17 together and with inclusion of the nitrogen atom to which they are bonded, form a 4-morpholinyl-, 1-pyrrolidinyl-, 1-piperidinyl-, 1-hexahydroazepino- or a 1-piperazinyl-ring of formula (c)




wherein

- R21 is 1-4C-alkyl, pyrid-4-yl, pyrid-4-ylmethyl, dimethylamino-1-4C-alkyl, dimethylaminocarbonylmethyl, N-methyl-piperidin-4-yl, 4-morpholino-ethyl or tetrahydrofuran-2-ylmethyl,
- R18 is halogen, nitro, cyano, carboxyl, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, amino, mono- or di-1-4C-alkylamino, aminocarbonyl, 1-4C-alkylcarbonylamino or mono- or di-1-4C-alkylaminocarbonyl,
- R19 is halogen, amino, nitro, 1-4C-alkyl or 1-4C-alkoxy,
- R20 is halogen,
- Aryl1 is pyrimidin-2-yl, thieno-[2,3-d]pyrimidin-4-yl, 1-methyl-1H-pyrazolo-[3,4-d]pyrimidin-4-yl, thiazolyl, imidazolyl, furanyl, pyridyl, phenyl or phenyl substituted by R18 and/or R19,
- Aryl2 is pyridyl, phenyl, phenyl substituted by R18 and/or R19, 2-oxo-2H-chromen-7-yl or 4-(1,2,3-thiadiazol-4-yl)phenyl,
- n is an integer from 1 to 4,
- m is an integer from 1 to 4,
- and the salts of these compounds.

Compounds of formula 1 of embodiment C to be emphasized are those in which

- R1 is hydrogen,
- R2 is hydrogen or 1-4C-alkyl,
- R3 represents a phenyl derivative of formulae (a) or (b)





wherein

- R4 is 1-2C-alkoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,
 - R5 is 1-4C-alkoxy,
 - R6 is 1-2C-alkoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,
 - R7 is methyl and
 - R8 is hydrogen,
 - or wherein
 - R7 and R8 together and with inclusion of the two carbon atoms, to which they are bonded, form a spiro-linked cyclopentane, cyclohexane, tetrahydrofurane or tetrahydropyran ring,
 - R9 is hydrogen, 1-4C-alkyl, $-\text{S}(\text{O})_2\text{-R10}$, $-\text{C}(\text{O})\text{R13}$, $-(\text{CH}_2)_m\text{-C}(\text{O})\text{-R15}$, Aryl1 or (Aryl2)-1-4C-alkyl,
 - R10 is 1-4C-alkyl, 5-dimethylaminonaphthalin-1-yl or $-\text{N}(\text{R16})\text{R17}$,
 - R13 is 1-4C-alkyl, phenyl, 4-ethyl-piperazin-2,3-dion-1-yl or $-\text{N}(\text{R16})\text{R17}$,
 - R15 is $-\text{N}(\text{R16})\text{R17}$, phenyl or phenyl substituted by R18 and/or R19 and/or R20,
 - R16 and R17 are independent from each other hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-methyl, phenyl or phenyl substituted by R18 and/or R19 and/or R20, or R16 and R17 together and with inclusion of the nitrogen atom to which they are bonded, form a 4-morpholinyl-, 1-pyrrolidinyl-, 1-piperidinyl-, 1-hexahydroazepino- or a 4-methyl-piperazin-1-yl-ring,
 - R18 is halogen, nitro, cyano, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxycarbonyl,
 - R19 is halogen, amino, nitro, 1-4C-alkyl or 1-4C-alkoxy,
 - R20 is halogen,
 - Aryl1 is pyrimidin-2-yl, thieno-[2,3-d]pyrimidin-4-yl, 1-methyl-1H-pyrazolo-[3,4-d]pyrimidin-4-yl, pyridyl, phenyl or phenyl substituted by R18 and/or R19,
 - Aryl2 is pyridyl, phenyl, phenyl substituted by R18 and/or R19, 2-oxo-2H-chromen-7-yl or 4-(1,2,3-thiadiazol-4-yl)phenyl,
 - m is an integer from 1 to 2,
- and the salts of these compounds.

Compounds of formula 1 of embodiment C particularly to be emphasized are those in which

- R1 is hydrogen,
- R2 is hydrogen or methyl,
- R3 represents a phenyl derivative of formulae (a) or (b)



wherein

R4 is 1-2C-alkoxy,

R5 is 1-4C-alkoxy,

R6 is 1-2C-alkoxy,

R7 is methyl and

R8 is hydrogen,

R9 is hydrogen, $-S(O)_2-R_{10}$, $-C(O)R_{13}$, $-(CH_2)_m-C(O)-R_{15}$ or (Aryl₂)-1-2C-alkyl,

R₁₀ is 1-4C-alkyl, 5-dimethylaminonaphthalin-1-yl or $-N(R_{16})R_{17}$,

R₁₃ is 1-4C-alkyl, 4-ethyl-piperazin-2,3-dion-1-yl or $-N(R_{16})R_{17}$,

R₁₅ is $-N(R_{16})R_{17}$,

R₁₆ and R₁₇ are independent from each other hydrogen or 1-4C-alkyl, or R₁₆ and R₁₇ together and with inclusion of the nitrogen atom to which they are bonded, form a 4-morpholinyl ring, a 1-piperidinyl ring or a 4-methyl-piperazin-1-yl ring,

Aryl₂ is pyridyl or phenyl,

m is 1,


and the salts of these compounds.

Preferred compounds of formula 1 of embodiment C are those in which

R₁ is hydrogen,

R₂ is methyl,

R₃ represents a phenyl derivative of formula (a)



wherein

R4 is methoxy,

R5 is methoxy and


R9 is acetyl, morpholin-4-ylcarbonyl, pyridin-3-ylmethyl, 4-ethyl-piperazin-2,3-dion-1-yl, 4-methylpiperazin-1-yl, 5-dimethylamino-naphthalene-1-sulfonyl or morpholin-4-yl-2-oxo-ethyl, and the salts of these compounds.

A special embodiment of the compounds of the present invention include those compounds of formula 1 in which R3 represents a phenyl derivative of formula (a).

Another special embodiment of the compounds of the present invention include those compounds of formula 1 in which R3 represents a phenyl derivative of formula (a) and R4 and R5 have the meaning methoxy.

A further special embodiment of the compounds of the present invention include those compounds of formula 1 in which R1 is hydrogen, R2 is hydrogen or methyl, R3 represents a phenyl derivative of formula (a) and R4 and R5 have the meaning methoxy.


The compounds of formula 1 are can be chiral compounds. Chiral centers exist in the compounds of formula 1 in positions 4 and 5 of the pyridazinone ring, if R1 and/or R2 have a meaning other than hydrogen. In case R3 represents a phenyl derivative of formula (b) there is one further chiral center in the dihydrofuran-ring, if the substituents -R7 and -CH₂R8 are not identical. However, preferred are in this connection those compounds, in which the substituents -R7 and -CH₂R8 are identical or together and with inclusion of the two carbon atoms to which they are bonded form a spiro-connected 5-, 6- or 7-membered hydrocarbon ring.



The invention includes all conceivable pure diastereomers and pure enantiomers of the compounds of formula 1, as well as all mixtures thereof independent from the ratio, including the racemates.

The compounds according to the invention can be prepared, for example, as described in Reaction scheme 1.

Reaction scheme 1:



Reaction scheme 1 shows that the compounds of formula 1 can be, for example, prepared starting from 4-oxo-piperidine-1-carboxylic acid tert-butyl ester which is reacted in a first reaction step with tert-butylcarbazate to give 4-(tert-Butoxycarbonyl-hydrazono)-piperidine-1-carboxylic acid tert-butyl ester (starting compound A6). Compound A6 is reduced with, for example, the boran tetrahydrofuran complex to give 4-(N'-tert-Butoxycarbonyl-hydrazino)-piperidine-1-carboxylic acid tert-butyl ester (starting

compound A5). Treatment of compound A5 with concentrated hydrochloric acid results in the formation of piperidin-4-yl-hydrazine dihydrochloride (starting compound A1).

The reaction of piperidin-4-yl-hydrazine dihydrochloride with phenyl-4-oxo-butyric acids of formulae 3a or 3b leads to the piperidino derivatives of formula 2.

These are reacted in the final reaction step with compounds of formula R9-X, wherein X represents a suitable leaving group, preferably a chlorine atom, to give the compounds of formula 1.

Suitably, the conversions are carried out analogous to methods which are familiar per se to the person skilled in the art, for example, in the manner which is described in the following examples.

The preparation of the phenyl-4-oxo-butyric acids of formulae 3a or 3b is known to the person skilled in the art (see for example Starting compounds and Intermediates).

The preparation of compounds of formula R9-X is also known to the person skilled in the art.

The substances according to the invention are isolated and purified in a manner known per se, e.g. by distilling off the solvent in vacuo and recrystallising the residue obtained from a suitable solvent or subjecting it to one of the customary purification methods, such as column chromatography on a suitable support material.

Salts are obtained by dissolving the free compound in a suitable solvent (for example a ketone like acetone, methylethylketone, or methylisobutylketone, an ether, like diethyl ether, tetrahydrofuran or dioxane, a chlorinated hydrocarbon, such as methylene chloride or chloroform, or a low molecular weight aliphatic alcohol, such as ethanol, isopropanol) which contains the desired acid, or to which the desired acid is then added. The salts are obtained by filtering, reprecipitating, precipitating with a non-solvent for the addition salt or by evaporating the solvent. Salts obtained can be converted by basification into the free compounds which, in turn, can be converted into salts. In this manner, pharmacologically non-tolerable salts can be converted into pharmacologically tolerable salts.

The following examples illustrate the invention in greater detail, without restricting it. As well, further compounds of formula 1, of which the preparation is explicitly not described, can be prepared in an analogous way or in a way which is known by a person skilled in the art using customary preparation methods.

The compounds, which are mentioned in the examples as well as their salts are preferred compounds of the invention. In the examples, RT stands for room temperature, h for hour(s), min for minute(s) and M. p. for melting point.

ExamplesFinal products

1. 6-(3,4-Dimethoxy-phenyl)-5-methyl-2-piperidin-4-yl-4,5-dihydro-2H-pyridazin-3-one hydrochloride

A mixture of 50 mmol of starting compound A1, 50 mmol of starting compound A2 and 100 mmol of triethylamine in 100 ml of 1-propanol is refluxed for 18 h and subsequently evaporated. The residue is partitioned between dichloromethane and aqueous sodium carbonate. The dichloromethane solution is dried over magnesium sulfate. Addition of a saturated solution of hydrochloric acid in diethyl ether causes precipitation of the title compound. M. p. 91-95°C

2. 6-(3,4-Dimethoxy-phenyl)-2-piperidin-4-yl-4,5-dihydro-2H-pyridazin-3-one hydrochloride

Prepared as described for compound 1 from starting compounds A1 and A3. M. p. 227-229°C

3. 6-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-2-piperidin-4-yl-4,5-dihydro-2H-pyridazin-3-one hydrochloride

Prepared as described for compound 1 from starting compounds A1 and A4. M. p. 280°C (with decomposition)

4. 2-(1-Acetyl-piperidin-4-yl)-6-(3,4-dimethoxy-phenyl)-5-methyl-4,5-dihydro-2H-pyridazin-3-one

To a solution of 5 mmol of compound 1 and 20 mmol of triethylamine in 50 ml of dichloromethane, 10 mmol of acetic anhydride is added and the resulting mixture is stirred at RT. After 60 min the solution is washed subsequently with diluted hydrochloric acid and aqueous sodium carbonate. The solution is dried over magnesium sulfate and evaporated. The residue is crystallized from diethyl ether. M. p. 149-152°C

5. 6-(3,4-Dimethoxy-phenyl)-5-methyl-2-[1-(1-morpholin-4-yl-methanoyl)-piperidin-4-yl]-4,5-dihydro-2H-pyridazin-3-one

Prepared from compound 1 and morpholine-4-carbonyl chloride as described for compound 4. M. p. 137-138°C

6. 6-(3,4-Dimethoxy-phenyl)-5-methyl-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4,5-dihydro-2H-pyridazin-3-one dihydrochloride

A mixture of 5 mmol of compound 1, 7 mmol of 3-chloromethyl-pyridine hydrochloride and 20 mmol of potassium carbonate in 20 ml of dimethylformamide is stirred at RT for 18 h. After addition of 150 ml of water, the mixture is extracted with diethyl ether. The ether solution is dried over magnesium sulfate and evaporated. The residue is purified by chromatography (elution with a mixture of ethyl acetate and methanol, 3:1). Addition of a saturated solution of hydrochloric acid in ether to the purified fractions causes precipitation of the title compound. M. p. 241-244°C

7. 1-(1-{4-[3-(3,4-Dimethoxy-phenyl)-6-oxo-5,6-dihydro-4H-pyridazin-1-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione

Prepared from compound 2 and 4-ethyl-2,3-dioxo-piperazine-1-carbonyl chloride as described for compound 4. M. p. 201-203°C

8. 6-(3,4-Dimethoxy-phenyl)-2-{1-[1-(4-methyl-piperazin-1-yl)-methanoyl]-piperidin-4-yl}-4,5-dihydro-2H-pyridazin-3-one hydrochloride

Prepared from compound 2 and 4-methyl-piperazine-1-carbonyl chloride as described for compound 4. After evaporating the dichloromethane solution, the residue is dissolved in ethyl acetate and addition of a saturated solution of hydrochloric acid in ether causes precipitation of the title compound. Recrystallisation is performed from a mixture of methanol and ethyl acetate. M. p. 151-154°C

9. 6-(3,4-Dimethoxy-phenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4,5-dihydro-2H-pyridazin-3-one

Prepared from compound 2 and 5-dimethylamino-naphthalene-1-sulfonyl chloride as described for compound 4. M. p. 191-193°C

10. 6-(3,4-Dimethoxy-phenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4,5-dihydro-2H-pyridazin-3-one hydrochloride

Prepared from compound 2 and 2-chloro-1-morpholin-4-yl-ethanone as described for compound 6. M. p. 145-148°C

11. 6-(3,4-Dimethoxy-phenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4,5-dihydro-2H-pyridazin-3-one

Prepared from 4-toluenesulfonyl chloride and compound 1 as described for compound 4.

M. p. 179-184°C

12. 6-(3,4-Dimethoxy-phenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4,5-dihydro-2H-pyridazin-3-one

Prepared from methylsulfonyl chloride and compound 1 as described for compound 4. M. p. 164-166°C

13. 2-[1-(4-Chloro-benzenesulfonyl)-piperidin-4-yl]-6-(3,4-dimethoxy-phenyl)-4,5-dihydro-2H-pyridazin-3-one

Prepared from 4-chloro-benzenesulfonyl chloride and compound 1 as described for compound 4.
M. p. 185-186°C

14. 6-(3,4-Dimethoxy-phenyl)-2-(1-phenylmethanesulfonyl-piperidin-4-yl)-4,5-dihydro-2H-pyridazin-3-one

Prepared from phenylmethanesulfonyl chloride and compound 1 as described for compound 4.
M. p. 114°C

15. 6-(3,4-Dimethoxy-phenyl)-2-[1-(4-methoxy-benzenesulfonyl)-piperidin-4-yl]-4,5-dihydro-2H-pyridazin-3-one

Prepared from 4-methoxy-benzenesulfonyl chloride and compound 1 as described for compound 4.
M. p. 197-198°C

16. 2-(1-Benzenesulfonyl-piperidin-4-yl)-6-(3,4-dimethoxy-phenyl)-4,5-dihydro-2H-pyridazin-3-one

Prepared from benzenesulfonyl chloride and compound 1 as described for compound 4.
M. p. 188-190°C

17. 2-[1-(2,5-Dimethoxy-benzenesulfonyl)-piperidin-4-yl]-6-(3,4-dimethoxy-phenyl)-4,5-dihydro-2H-pyridazin-3-one

Prepared from 2,5-Dimethoxy-benzenesulfonyl chloride and compound 1 as described for compound 4. M. p. 184-185°C

18. 2-[4-[3-(3,4-Dimethoxy-phenyl)-6-oxo-5,6-dihydro-4H-pyridazin-1-yl]-piperidine-1-sulfonyl]-benzonitrile

Prepared from 2-cyano-benzenesulfonyl chloride and compound 1 as described for compound 4.

M. p. 158-160 °C

19. 6-(3,4-Dimethoxy-phenyl)-2-[1-(thiophene-2-sulfonyl)-piperidin-4-yl]-4,5-dihydro-2H-pyridazin-3-one

Prepared from 2-thiophene sulfonyl chloride and compound 1 as described for compound 4.

M. p. 178-179 °C

20. 6-(3,4-Dimethoxy-phenyl)-2-[1-(2-fluoro-benzenesulfonyl)-piperidin-4-yl]-4,5-dihydro-2H-pyridazin-3-one

Prepared from 2-fluoro-benzenesulfonyl chloride and compound 1 as described for compound 4.

M. p. 198-199 °C

21. 6-(3,4-Dimethoxy-phenyl)-2-[1-(2-trifluoromethoxy-benzenesulfonyl)-piperidin-4-yl]-4,5-dihydro-2H-pyridazin-3-one

Prepared from 2-trifluoromethoxy-benzenesulfonyl chloride and compound 1 as described for compound 4. M. p. 118-119 °C

22. 4-[3-(3,4-Dimethoxy-phenyl)-6-oxo-5,6-dihydro-4H-pyridazin-1-yl]-piperidine-1-sulfonic acid dimethylamide

Prepared from dimethylsulfamoyl chloride and compound 1 as described for compound 4.

M. p. 103-106 °C

23. 6-(3,4-Dimethoxy-phenyl)-2-[1-(1-phenyl-methanoyl)-piperidin-4-yl]-4,5-dihydro-2H-pyridazin-3-one

Prepared from benzoyl chloride and compound 1 as described for compound 4.

M. p. 152-154 °C

24. 6-(3,4-Dimethoxy-phenyl)-2-[1-(1-pyridin-3-yl-methanoyl)-piperidin-4-yl]-4,5-dihydro-2H-pyridazin-3-one

Prepared from nicotinoyl chloride and compound 1 as described for compound 4.

M. p. 162-164 °C

25. 6-(3,4-Dimethoxy-phenyl)-2-{1-[1-(2,4,6-trichloro-phenyl)-methanoyl]-piperidin-4-yl}-4,5-dihydro-2H-pyridazin-3-one

Prepared from 2,4,6-trichlorobenzoyl chloride and compound 1 as described for compound 4.

M. p. 156-159 °C

26. 4-[3-(3,4-Dimethoxy-phenyl)-6-oxo-5,6-dihydro-4H-pyridazin-1-yl]-piperidine-1-carboxylic acid tert-butylamide

Prepared from tert-butyl isocyanate and compound 1 as described for compound 4.

M. p. 76-78 °C

27. 2-[4-[3-(3,4-Dimethoxy-phenyl)-6-oxo-5,6-dihydro-4H-pyridazin-1-yl]-piperidin-1-yl]-N,N-dimethylacetamide hydrochloride

Prepared from 2-chloro-N,N-dimethylacetamide and compound 1 as described for compound 6.

M. p. 121-122 °C

28. 6-(3,4-Dimethoxy-phenyl)-2-{1-[2-(4-methyl-piperazin-1-yl)-2-oxo-ethyl]-piperidin-4-yl}-4,5-dihydro-2H-pyridazin-3-one dihydrochloride

Prepared from 2-chloro-1-(4-methyl-piperazin-1-yl)-ethanone hydrochloride and compound 1 as described for compound 6. M. p. 194-199 °C

29. 6-(3,4-Dimethoxy-phenyl)-2-{1-[2-(4-pyridin-4-yl-piperazin-1-yl)-ethanoyl]-piperidin-4-yl}-4,5-dihydro-2H-pyridazin-3-one dihydrochloride

Prepared from 1-Pyridin-4-yl-piperazine and starting compound A7 as described for compound 6.

M. p. 98-99 °C

30. 6-(3,4-Dimethoxy-phenyl)-2-(1-{2-[4-(2-methoxy-phenyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4,5-dihydro-2H-pyridazin-3-one dihydrochloride

Prepared from 1-(2-methoxy-phenyl)-piperazine and compound A7 as described for compound 6.

M. p. 109-110 °C

31. 6-(3,4-Dimethoxy-phenyl)-2-(1-{2-[4-(1,1-diphenyl-methyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4,5-dihydro-2H-pyridazin-3-one dihydrochloride

Prepared from 1-(1,1-diphenyl-methyl)-piperazine and compound A7 as described for compound 6.

M. p. 126-127 °C

Starting Compounds and Intermediates**A1. Piperidin-4-yl-hydrazine dihydrochloride**

A mixture of 0.1 mole of 4-(N'-tert-Butoxycarbonyl-hydrazino)-piperidine-1-carboxylic acid tert-butyl ester (starting compound A5) and 150 ml of concentrated hydrochloric acid is heated at 90°C for 60 min after which the clear solution is evaporated. The residue is washed with tetrahydrofurane, filtered off and dried under vacuum. M. p. 256-259°C

A2. 4-(3,4-Dimethoxy-phenyl)-3-methyl-4-oxo-butyric acid

Prepared according to Haworth and Woodcock; J. Chem. Soc. 1938, 809-811

A3. 4-(3,4-Dimethoxy-phenyl)-4-oxo-butyric acid

Prepared according to M.S.Y. Khan and Anees A. Siddiqui; Indian J. Chem. Section B, 2000, 39, 614-619

A4. 4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-4-oxo-butyric acid

Prepared analogously to (cis)-2-(2,3-Dihydro-7-methoxybenzofuran-2-spiro-1'-cyclopentane-4-carbonyl)-1,2,3,6-tetrahydrobenzoic acid as described in WO99/31090 starting from 4-bromo-7-methoxy-2,2-dimethyl-2,3-dihydro-benzofuran and succinic anhydride. M. p. 125-126°C

A5. 4-(N'-tert-Butoxycarbonyl-hydrazino)-piperidine-1-carboxylic acid tert-butyl ester

150 ml of a solution of borohydride in tetrahydrofurane (1.0 mol/l) is slowly added to a solution of 0.12 mole of 4-(tert-Butoxycarbonyl-hydrazono)-piperidine-1-carboxylic acid tert-butyl ester (starting compound A6) in 100 ml of dry tetrahydrofurane. After complete addition, the mixture is stirred for another 30 min after which a 100 ml of water is added to destroy the excess of borohydride. Subsequently the tetrahydrofurane is evaporated and the resulting aqueous solution is extracted with diethyl ether. After drying the solvent over magnesium sulfate, the ether is evaporated. M. p. 112-115°C

A6. 4-(tert-Butoxycarbonyl-hydrazono)-piperidine-1-carboxylic acid tert-butyl ester

A mixture of 0.15 mole of 4-oxo-piperidine-1-carboxylic acid tert-butyl ester and 0.15 mole of tert-butylcarbazate in 250 ml of hexane is stirred for 18 h at RT. The precipitate is filtered off and dried under vacuum. M. p. 172-174°C

A7. 2-[1-(2-Chloro-ethanoyl)-piperidin-4-yl]-6-(3,4-dimethoxy-phenyl)-4,5-dihydro-2H-pyridazin-3-one

A solution of 20 mmol of chloroacetyl chloride in 50 ml of dichloromethane is added slowly to a solution of 15 mmol of compound 1 and 40 mmol of triethylamine in 150 ml of dichloromethane at 0°C. After complete addition, the mixture is washed successively with diluted hydrochloric acid and with aqueous sodium carbonate, dried over magnesium sulfate and evaporated. The compound is crystallized from ethyl acetate. M. p. 116-117 °C

Commercial utility

The compounds according to the invention have valuable pharmacological properties which make them commercially utilizable. As selective inhibitors of type 4 or type 3 and 4 of cyclic nucleotide phosphodiesterase (PDE4, PDE3/4), they are suitable on the one hand as bronchial therapeutics (for the treatment of airway obstructions on account of their dilating action and cilia-stimulating action but also on account of their respiratory rate- and respiratory drive-increasing action), but on the other hand especially for the treatment of disorders of inflammatory nature, e.g. of the airways (asthma prophylaxis), of the skin, of the intestine, of the eyes and of the joints, which are mediated by mediators such as interferons, members of the tumour necrosis factor family, interleukins, chemokines, colony-stimulating factors, growth factors, lipid mediators (e.g., inter alia, PAF, platelet-activating factor), bacterial factors (e.g. LPS), immunoglobulins, oxygen free radicals and related free radicals (e.g. nitrogen monoxide NO), biogenic amines (e.g. histamine, serotonin), kinins (e.g. bradykinin), neurogenic mediators (such as substance P, neurokinin), proteins such as, for example, granular contents of leukocytes (inter alia cationic proteins of eosinophils) and adherence proteins (e.g. integrins). The compounds according to the invention have smooth muscle-relaxant action, e.g. in the region of the bronchial system, of the blood circulation, and of the efferent urinary passages. Furthermore, they have cilia frequency-increasing action, for example in the bronchial system.

On account of their PDE-inhibiting properties, the compounds according to the invention can be employed as therapeutics in human and veterinary medicine, where they can be used, for example, for the treatment and prophylaxis of the following diseases: acute and chronic (in particular inflammatory and allergen-induced) respiratory disorders of various origins (bronchitis, allergic bronchitis, bronchial asthma, emphysema, COPD); disorders associated with impaired cilia function or increased demands on ciliar clearance (bronchitis, mucoviscidosis), dermatoses (especially of proliferative, inflammatory and allergic type) such as, for example, psoriasis (vulgaris), toxic and allergic contact eczema, atopic eczema, seborrheic eczema, lichen simplex, sunburn, pruritus in the anogenital area, alopecia areata, hypertrophic scars, discoid lupus erythematosus, follicular and widespread pyodermites, endogenous and exogenous acne, acne rosacea and other proliferative, inflammatory and allergic skin disorders; disorders which are based on excessive release of TNF and leukotrienes, i.e., for example, disorders of the arthritis type (rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis and other arthritic conditions), systemic lupus erythematosus, disorders of the immune system (AIDS), including AIDS-related encephalopathies, autoimmune disorders such as diabetes mellitus (type I, autoimmune diabetes), multiple sclerosis and of the type virus-, bacteria- or parasite-induced demyelination diseases, cerebral malaria or Lyme's disease, shock symptoms [septic shock, endotoxin shock, Gram-negative sepsis, toxic shock syndrome and ARDS (adult respiratory distress syndrome)] and also generalized inflammations in the gastrointestinal region (Crohn's disease and ulcerative colitis); disorders which are based on allergic and/or chronic, faulty immunological reactions in the region of the upper airways (pharynx, nose) and of the adjacent regions (paranasal sinuses, eyes), such as, for example, allergic rhinitis/sinusitis, chronic rhinitis/sinusitis, allergic conjunctivitis and also nasal polyps; and also disorders

of the central nervous system such as memory disorders and Alzheimer's disease, candidiasis, leishmaniasis and leprosy.

On account of their vasorelaxant activity, the compounds according to the invention can also be used for the treatment of high blood pressure disorders of various origins such as, for example, pulmonary high blood pressure and the concomitant symptoms associated therewith, for the treatment of erectile dysfunction or colics of the kidneys and the ureters in connection with kidney stones.

On account of their cAMP-increasing action, however, they can also be used for disorders of the heart which can be treated by PDE inhibitors, such as, for example, cardiac insufficiency, and also as anti-thrombotic, platelet aggregation-inhibiting substances.

The invention further relates to a method for the treatment of mammals including humans who are suffering from one of the abovementioned diseases. The method comprises administering a therapeutically effective and pharmacologically acceptable amount of one or more of the compounds according to the invention to the sick mammal.

The invention further relates to the compounds according to the invention for use in the treatment and/or prophylaxis of diseases, in particular the diseases mentioned.

The invention also relates to the use of the compounds according to the invention for the production of pharmaceutical compositions which are employed for the treatment and/or prophylaxis of the diseases mentioned.

The invention furthermore relates to pharmaceutical compositions for the treatment and/or prophylaxis of the diseases mentioned and which contain one or more of the compounds according to the invention.

A further subject of the invention is a commercial product, consisting of a customary secondary pack, a primary pack containing the pharmaceutical composition (for example an ampoule or a blister pack) and, if desired, an information leaflet, the pharmaceutical composition exhibiting antagonistic action against cyclic nucleotide phosphodiesterases of type 4 or types 3 and 4 and leading to the attenuation of the symptoms of illnesses which are connected with cyclic nucleotide phosphodiesterases of type 4 or types 3 and 4, and the suitability of the pharmaceutical composition for the prophylaxis or treatment of illnesses which are connected with cyclic nucleotide phosphodiesterases of type 4 or types 3 and 4 being indicated on the secondary pack and/or on the information leaflet of the commercial product, and the pharmaceutical composition containing one or more compounds of formula 1 according to the invention. The secondary pack, the primary pack containing the pharmaceutical composition and the information leaflet otherwise comply with what would be regarded as standard to the person skilled in the art for pharmaceutical compositions of this type.

Advantageously, the substances according to the invention are also suitable for combination with other substances which bring about stimulation of cAMP, such as prostaglandins (PGE₂, PGI₂ and prostacyclin) and their derivatives, direct adenylate cyclase stimulators such as forskolin and related substances, or substances indirectly stimulating adenylate cyclase, such as catecholamines and adrenergic receptor agonists, in particular beta-mimetics. In combination, on account of their cAMP degradation-inhibiting action, they in this case display a synergistic, superadditive activity. This comes to bear, for example, in their use in combination with PGE₂ for the treatment of pulmonary hypertension.

The pharmaceutical compositions are prepared by processes which are known per se and familiar to the person skilled in the art. As pharmaceutical compositions, the compounds according to the invention (= active compounds) are either employed as such, or preferably in combination with suitable pharmaceutical auxiliaries and/or excipients, e.g. in the form of tablets, coated tablets, capsules, caplets, suppositories, patches (e.g. as TTS), emulsions, suspensions, gels or solutions, the active compound content advantageously being between 0.1 and 95% and where, by the appropriate choice of the auxiliaries and/or excipients, a pharmaceutical administration form (e.g. a delayed release form or an enteric form) exactly suited to the active compound and/or to the desired onset of action can be achieved.

The person skilled in the art is familiar with auxiliaries or excipients which are suitable for the desired pharmaceutical formulations on account of his/her expert knowledge. In addition to solvents, gel formers, ointment bases and other active compound excipients, for example antioxidants, dispersants, emulsifiers, preservatives, solubilizers, colorants, complexing agents or permeation promoters, can be used.

The administration of the pharmaceutical compositions according to the invention may be performed in any of the generally accepted modes of administration available in the art. Illustrative examples of suitable modes of administration include intravenous, oral, nasal, parenteral, topical, transdermal and rectal delivery. Oral delivery is preferred.

For the treatment of disorders of the respiratory tract, the compounds according to the invention are preferably also administered by inhalation in the form of an aerosol; the aerosol particles of solid, liquid or mixed composition preferably having a diameter of 0.5 to 10 µm, advantageously of 2 to 6 µm.

Aerosol generation can be carried out, for example, by pressure-driven jet atomizers or ultrasonic atomizers, but advantageously by propellant-driven metered aerosols or propellant-free administration of micronized active compounds from inhalation capsules.

Depending on the inhaler system used, in addition to the active compounds the administration forms additionally contain the required excipients, such as, for example, propellants (e.g. Frigen in the case of

metered aerosols), surface-active substances, emulsifiers, stabilizers, preservatives, flavorings, fillers (e.g. lactose in the case of powder inhalers) or, if appropriate, further active compounds.

For the purposes of inhalation, a large number of apparatuses are available with which aerosols of optimum particle size can be generated and administered, using an inhalation technique which is as right as possible for the patient. In addition to the use of adaptors (spacers, expanders) and pear-shaped containers (e.g. Nebulator®, Volumatic®), and automatic devices emitting a puffer spray (Autohaler®), for metered aerosols, in particular in the case of powder inhalers, a number of technical solutions are available (e.g. Diskhaler®, Rotadisk®, Turbohaler® or the inhaler described in European Patent Application EP 0 505 321), using which an optimal administration of active compound can be achieved.

For the treatment of dermatoses, the compounds according to the invention are in particular administered in the form of those pharmaceutical compositions which are suitable for topical application. For the production of the pharmaceutical compositions, the compounds according to the invention (= active compounds) are preferably mixed with suitable pharmaceutical auxiliaries and further processed to give suitable pharmaceutical formulations. Suitable pharmaceutical formulations are, for example, powders, emulsions, suspensions, sprays, oils, ointments, fatty ointments, creams, pastes, gels or solutions.

The pharmaceutical compositions according to the invention are prepared by methods known per se. The dosage of the active compounds takes place in the order of magnitude customary for PDE inhibitors. Thus topical application forms (such as, for example, ointments) for the treatment of dermatoses contain the active compounds in a concentration of, for example, 0.1-99%. The dose for administration by inhalation is customarily between 0.1 and 3 mg per day. The customary dose in the case of systemic therapy (p.o. or i.v.) is between 0.01 and 10 mg per kilogram per day.

Biological investigations

The second messenger cyclic AMP (cAMP) is known for inhibiting inflammatory cells and cells responsible for the immunological response. The PDE4 isoenzyme is widely distributed in cells associated with the initiation and spreading of inflammatory diseases (H Tenor and C Schudt, in "Phosphodiesterase Inhibitors", 21-40, "The Handbook of Immunopharmacology", Academic Press 1996); its inhibition results in the increase of the intracellular cyclic AMP concentration and thus in the inhibition of cellular activation (JE Souness et al., Immunopharmacology 47: 127-162, 2000).

The anti-inflammatory potential of PDE4 inhibitors in vivo has been described in various animal models (MMTeixeira, TIPS 18: 164-170, 1997). To examine the PDE4 inhibition on a cellular level (in vitro), a large number of proinflammatory responses can be measured. Examples are the superoxide production of neutrophilic (C Schudt et al., Arch Pharmacol 344: 682-690, 1991) or eosinophilic (A Hatzelmann et al., Brit J Pharmacol 114: 821-831, 1995) granulocytes, which can be measured as luminol-enhanced chemiluminescence, or the synthesis of tumor necrosis factor alpha (TNF α) in monocytes, macrophages or dendritic cells (Gantner et al., Brit J Pharmacol 121: 221-231, 1997 and Pulmonary Pharmacol Therap 12: 377-386, 1999). The immunomodulatory potential of the PDE4 inhibitors furthermore becomes apparent by inhibition of T-cell responses such as cytokine synthesis or proliferation (DM Essayan, Biochem Pharmacol 57: 965-973, 1999). PDE4 inhibition by the substances according to the invention is thus a central indicator of the suppression of inflammatory processes.

Some of the cells involved in inflammatory processes contain, in addition to PDE4, also the PDE3 isoenzyme which likewise contributes to the total cAMP metabolism of these cells. Examples are endothelial cells, mast cells, T-cells, macrophages and dendritic cells. In these cell types, the inhibitory action of PDE4 inhibitors can be enhanced by additional PDE3 inhibition. In the case of (respiratory) smooth muscle cells, inhibition of the PDE3 activity is furthermore important for (broncho)relaxation (A Hatzelmann et al., in "Phosphodiesterase Inhibitors", 147-160, "The Handbook of Immunopharmacology", Academic Press, 1996).

Method for measuring inhibition of PDE3 and PDE4 activities**Method A:**

The PDE activity was determined according to Thompson et al. (Adv Cycl Nucl Res 10: 69-92, 1979) with some modifications (Bauer and Schwabe, Naunyn-Schmiedeberg's Arch Pharmacol 311: 193-198, 1980). The test samples contained 20 mM Tris (pH 7.4), 5 mM MgCl₂, 0.5 µM cAMP or cGMP, [³H]cAMP or [³H]cGMP (about 30 000 cpm/sample), the PDE isoenzyme-specific additives described in greater detail below, the indicated concentrations of inhibitor and an aliquot of the enzyme solution in a total sample volume of 200 µl. Dilution series of the compounds according to the invention were prepared in DMSO and further diluted in the samples [1:100 (v/v)], to give the desired end concentration of the inhibitors at a DMSO concentration of 1% (v/v), which for its part has only a minute effect on PDE activity.

After preincubation at 37°C for 5 minutes, the reaction was started by addition of the substrate (cAMP or cGMP). The samples were incubated at 37°C for a further 15 min. The reaction was terminated by addition of 50 µl 0.2 N HCl. After cooling on ice for 10 minutes and addition of 25 µg 5'-nucleotidase (snake venom from Crotalus atrox), the mixture was again incubated at 37°C for 10 min and the samples were then applied to QAE Sephadex A-25 columns (sample volume 1 ml). The columns were eluted with 2 ml of 30 mM ammonium formate (pH 6.0). The radioactivity of the eluate was measured and corrected by the corresponding blank values (measured in the presence of denatured protein); the blank values were less than 5% of the total radioactivity. In no case did the proportion of hydrolyzed nucleotide exceed 30% of the original substrate concentration.

PDE3 (cGMP-inhibited) was investigated in homogenates of human platelets (see Schudt et al., Biochem Pharmacol 1991: 42, 153-162) using cAMP or cGMP as substrate.

PDE4 (cAMP-specific) was investigated in the cytosol of human polymorphonuclear leukocytes (PMNL) [isolated from leukocyte concentrates, see Schudt et al., Arch Pharmacol 1991: 344, 682-690] using cAMP as substrate. The PDE3 inhibitor motapizone (1 µM) was used to suppress the PDE3 activity emanating from contaminated platelets.

The IC₅₀ values were determined from the concentration-inhibition curves by nonlinear regression.

Method B:

The cDNA for PDE3A1 (GB no. U36798) was isolated in 2 steps using PCR. A 3' terminal cDNA fragment of PDE3A1 was amplified from fat cells cDNA (Clontech, Palo Alto) using primers OZ 458 (5'-

AAAGTCGACTCACTGGTCTGGCTTTTGG -3') and OZ 457 (5'- GTCGACCAGGTGCCCTCGCTA - 3'). The 5' terminal cDNA fragment of PDE3A1 was amplified from Placenta cDNA (Clontech, Palo Alto) using primers OZ 455 (5'- ATGGCAGTGCCCGGCGACGCT -3') and OZ 456 (5'- GTCGACTTTGCTTTTTCAGCCT -3'). The PCR products were cloned into pCR2.1-Topo (Invitrogen, Groningen, NL) under standard conditions (the manufacturer's instructions). The 3' fragment was cut out with HindII and cloned into the HindII site of the construct carrying the 5' fragment. The whole ORF was subcloned into pBacPak9 (Clontech, Palo Alto) using EcoRI. Aminoacid 12 is Aspartic Acid like in sequence GB no. AJ005036, aa 69 and aa 110 are respective Serine and Glycine like in both sequences GB no. AJ005036 and GB no. M91667.

The PDE4B2 (GB no. M97515) was a gift of Prof. M. Conti (Stanford University, USA). It was amplified from the original plasmid (pCMV5) via PCR with primers Rb9 (5'- GCCAGCGTGCAAATAATGAAGG - 3') and Rb10 (5'- AGAGGGGGATTATGTATCCAC -3') and cloned into the pCR-Bac vector (Invitrogen, Groningen, NL).

The recombinant baculovirus was prepared by means of homologous recombination in SF9 insect cells. The expression plasmids were cotransfected with Bac-N-Blue (Invitrogen, Groningen, NL) or Baculo-Gold DNA (Pharmingen, Hamburg) using a standard protocol (Pharmingen, Hamburg). Wt virus-free recombinant virus supernatants were selected using plaque assay methods. After that, high-titre virus supernatants were prepared by amplifying 3 times. PDEs were expressed in SF21 cells by infecting 2×10^6 cells/ml with an MOI (multiplicity of infection) between 1 and 10 in serum-free SF900 medium (Life Technologies, Paisley, UK). The cells were cultured at 28°C for 48 – 72 hours, after which they were pelleted for 5-10 min at 1000 g and 4°C.

The SF21 insect cells were resuspended, at a concentration of approx. 10^7 cells/ml, in ice-cold (4°C) homogenization buffer (20 mM Tris, pH 8.2, containing the following additions: 140 mM NaCl, 3.8 mM KCl, 1 mM EGTA, 1 mM MgCl₂, 10 mM β -mercaptoethanol, 2 mM benzamidine, 0.4 mM Pefablock, 10 μ M leupeptin, 10 μ M pepstatin A, 5 μ M trypsin inhibitor) and disrupted by ultrasonication. The homogenate was then centrifuged for 10 min at 1000 \times g and the supernatant was stored at -80°C until subsequent use (see below). The protein content was determined by the Bradford method (BioRad, Munich) using BSA as the standard.

PDE3A1 and PDE4B2 activities were inhibited by the said compounds in a modified SPA (scintillation proximity assay) test, supplied by Amersham Biosciences (see procedural instructions "phosphodiesterase [3H]cAMP SPA enzyme assay, code TRKQ 7090"), carried out in 96-well microtitre plates (MTP's). The test volume is 100 μ l and contains 20 mM Tris buffer (pH 7.4), 0.1 mg of BSA (bovine serum albumin)/ml, 5 mM Mg²⁺, 0.5 μ M cAMP (including about 50,000 cpm of [3H]cAMP), 1 μ l of the respective substance dilution in DMSO and sufficient recombinant PDE (1000 \times g supernatant, see above) to ensure that 10-20% of the cAMP is converted under the said experimental conditions. The final concentration of DMSO in the assays (1 % v/v) does not substantially affect the activity of the PDEs investigated. After a preincubation of 5 min at 37°C, the reaction is started by adding the sub-

strate (cAMP) and the assays are incubated for a further 15 min; after that, they are stopped by adding SPA beads (50 μ l). In accordance with the manufacturer's instructions, the SPA beads had previously been resuspended in water, but were then diluted 1:3 (v/v) in water; the diluted solution also contains 3 mM IBMX to ensure a complete PDE activity stop. After the beads have been sedimented (> 30 min), the MTP's are analyzed in commercially available luminescence detection devices. The corresponding IC₅₀ values of the compounds for the inhibition of PDE activities are determined from the concentration-effect curves by means of non-linear regression.

The inhibitory values determined for the compounds according to the invention follow from the following Table 1, in which the numbers of the compounds correspond to the numbers of the examples.


The inhibitory values of the compounds 1-22 and 27 have been determined according to Method A. The inhibitory values of the compounds 23-26, 28 and 29-31 have been determined according to Method B.

Table 1**Inhibition of PDE4 and PDE3 activity [measured as $-\log IC_{50}$ (mol/l)]**

compound	PDE4 Inhibition	PDE3 Inhibition
4	8.00	5.76
5	7.89	5.75
6	8.39	5.23
7	8.96	< 5
8	7.71	< 5
9	7.53	6.61
10	7.17	5.35
11	8.3	6.3
12	7.8	6.6
14	8.2	6.6
15	7.8	6.3
17	7.8	6.8
18	8.0	7.1
19	8.3	6.9
20	8.4	6.8
21	8.5	7.0
23	8.1	6.9
25	8.6	7.8
27	9.2	< 5
28	7.7	< 5
29	7.8	5.5
31	7.4	5.4

Patent claims

1. Compounds of formula 1





in which

R1 is hydrogen or 1-4C-alkyl,

R2 is hydrogen or 1-4C-alkyl,

R3 represents a phenyl derivative of formulae (a) or (b)



wherein

R4 is 1-4C-alkoxy or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R5 is 1-8C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy, or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R6 is 1-4C-alkoxy, 3-5C-cycloalkoxy, 3-5C-cycloalkylmethoxy, or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R7 is 1-4C-alkyl and

R8 is hydrogen or 1-4C-alkyl,

or wherein

R7 and R8 together and with inclusion of the two carbon atoms, to which they are bonded, form a spiro-linked 5-, 6- or 7-membered hydrocarbon ring, optionally interrupted by an oxygen or sulphur atom,

R9 is 1-4C-alkyl, $-S(O)_2-R_{10}$, $-S(O)_2-(CH_2)_n-R_{11}$, $-(CH_2)_m-S(O)_2-R_{12}$, $-C(O)R_{13}$, $-C(O)-(CH_2)_n-R_{14}$, $-(CH_2)_m-C(O)-R_{15}$, Aryl1 or (Aryl2)-1-4C-alkyl,

R10 is 1-4C-alkyl, 5-dimethylaminonaphthalin-1-yl, $-N(R_{16})R_{17}$, thiophenyl, phenyl or phenyl substituted by R18 and/or R19,

R11 is phenyl or -N(R16)R17,

R12 is -N(R16)R17,

R13 is 1-4C-alkyl, hydroxycarbonyl-1-4C-alkyl, phenyl, 2,4,6-trichlorophenyl, pyridyl, 4-ethyl-piperazin-2,3-dion-1-yl or -N(R16)R17,

R14 is -N(R16)R17,


R15 is -N(R16)R17, phenyl or phenyl substituted by R18 and/or R19 and/or R20,

R16 is hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkylmethyl, phenyl or phenyl substituted by R18 and/or R19 and/or R20,

R17 1-7C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkylmethyl, phenyl or phenyl substituted by R18 and/or R19 and/or R20,

or

R16 and R17 together and with inclusion of the nitrogen atom to which they are bonded, form a 4-morpholinyl-, 1-pyrrolidinyl-, 1-piperidinyl-, 1-hexahydroazepino- or a 1-piperazinyl-ring of formula (c)



wherein

R21 is 1-4C-alkyl, pyrid-4-yl, pyrid-4-ylmethyl, 2-methoxyphenyl, 1,1-diphenylmethyl, dimethyl-amino-1-4C-alkyl, dimethylaminocarbonylmethyl, N-methyl-piperidin-4-yl, 4-morpholino-ethyl or tetrahydrofuran-2-ylmethyl,

R18 is halogen, nitro, cyano, carboxyl, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine, 1-4C-alkoxycarbonyl, amino, mono- or di-1-4C-alkylamino, aminocarbonyl, 1-4C-alkylcarbonylamino or mono- or di-1-4C-alkylaminocarbonyl,

R19 is halogen, amino, nitro, 1-4C-alkyl or 1-4C-alkoxy,

R20 is halogen,

Aryl1 is pyrimidin-2-yl, thieno-[2,3-d]pyrimidin-4-yl, 1-methyl-1H-pyrazolo-[3,4-d]pyrimidin-4-yl, thiazolyl, imidazolyl, furanyl, pyridyl, phenyl or phenyl substituted by R18 and/or R19,

Aryl2 is pyridyl, phenyl, phenyl substituted by R18 and/or R19, 2-oxo-2H-chromen-7-yl or 4-(1,2,3-thiadiazol-4-yl)phenyl,

n is an integer from 1 to 4,

m is an integer from 1 to 4,



and the salts of these compounds.

2. Compounds of formula 1 according to claim 1 in which

R1 is hydrogen,


R2 is hydrogen or 1-4C-alkyl,

R3 represents a phenyl derivative of formulae (a) or (b)



wherein

- R4 is 1-2C-alkoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,
 R5 is 1-4C-alkoxy,
 R6 is 1-2C-alkoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,
 R7 is methyl and
 R8 is hydrogen,
 or wherein
 R7 and R8 together and with inclusion of the two carbon atoms, to which they are bonded, form a
 spiro-linked cyclopentane, cyclohexane, tetrahydrofurane or tetrahydropyran ring,
 R9 is 1-4C-alkyl, -S(O)₂-R10, -S(O)₂-(CH₂)_n-R11, -C(O)R13, -C(O)-(CH₂)_n-R14, -(CH₂)_m-C(O)-R15 or
 (Aryl)₂-1-4C-alkyl,
 R10 is 1-4C-alkyl, 5-dimethylaminonaphthalin-1-yl, -N(R16)R17, thiophenyl, phenyl or phenyl substi-
 tuted by R18 and/or R19,
 R11 is phenyl,
 R13 is 1-4C-alkyl, phenyl, pyridyl, 2,4,6-trichlorophenyl, 4-ethyl-piperazin-2,3-dion-1-yl or -N(R16)R17,
 R14 is -N(R16)R17,
 R15 is -N(R16)R17, phenyl or phenyl substituted by R18 and/or R19,
 R16 is hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl or 3-7C-cycloalkylmethyl,
 R17 is 1-7C-alkyl, 3-7C-cycloalkyl or 3-7C-cycloalkylmethyl,
 or
 R16 and R17 together and with inclusion of the nitrogen atom to which they are bonded, form a
 4-morpholinyl-, 1-pyrrolidinyl-, 1-piperidinyl-, 1-hexahydroazepino- or a 1-piperazinyl-ring of for-
 mula (c)



wherein

R21 is 1-4C-alkyl, pyrid-4-yl, 2-methoxyphenyl, 1,1-diphenylmethyl or N-methyl-piperidin-4-yl,


R18 is halogen, nitro, cyano, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxy which is completely or predominantly substituted by fluorine or 1-4C-alkoxycarbonyl,
 R19 is halogen, amino, nitro, 1-4C-alkyl or 1-4C-alkoxy,
 Aryl2 is pyridyl, phenyl or phenyl substituted by R18 and/or R19,
 m is an integer from 1 to 2,
 n is an integer from 1 to 2,
 and the salts of these compounds.

3. Compounds of formula 1 according to claim 1 in which

R1 is hydrogen,

R2 is hydrogen or methyl,

R3 represents a phenyl derivative of formula (a)



wherein

R4 is 1-2C-alkoxy,

R5 is 1-2C-alkoxy,

R9 is $-S(O)_2-R_{10}$, $-S(O)_2-(CH_2)_n-R_{11}$, $-C(O)R_{13}$, $-C(O)-(CH_2)_n-R_{14}$, $-(CH_2)_m-C(O)-R_{15}$ or (Aryl2)-1-2C-alkyl,

R10 is 1-4C-alkyl, 5-dimethylaminonaphthalin-1-yl, $-N(R_{16})R_{17}$, thiophenyl, phenyl or phenyl substituted by R18 and/or R19,

R11 is phenyl,

R13 is 1-4C-alkyl, phenyl, 2,4,6-trichlorophenyl, pyridyl, 4-ethyl-piperazin-2,3-dion-1-yl or $-N(R_{16})R_{17}$,

R14 is $-N(R_{16})R_{17}$,


R15 is $-N(R_{16})R_{17}$,

R16 is hydrogen or 1-4C-alkyl,

R17 is 1-4C-alkyl,

or


R16 and R17 together and with inclusion of the nitrogen atom to which they are bonded, form a 4-morpholinyl-, 1-pyrrolidinyl-, 1-piperidinyl- or a 1-piperazinyl-ring of formula (c)



wherein

R21 is 1-4C-alkyl, pyrid-4-yl, 2-methoxyphenyl or 1,1-diphenylmethyl,
 R18 is halogen, cyano, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,
 R19 is 1-4C-alkyl or 1-4C-alkoxy,
 Aryl2 is pyridyl or phenyl,
 m is 1,
 n is 1,
 and the salts of these compounds.


4. Compounds of formula 1 according to claim 1 in which
 R1 is hydrogen,
 R2 is hydrogen or methyl,
 R3 represents a phenyl derivative of formula (a)



wherein

R4 is methoxy,
 R5 is methoxy and
 R9 is acetyl, morpholin-4-ylcarbonyl, pyridin-3-ylmethyl, 4-ethyl-piperazin-2,3-dione-1-ylcarbonyl, 4-methylpiperazin-1-ylcarbonyl, 5-dimethylamino-naphthalene-1-sulfonyl, 2-(morpholin-4-yl)-2-oxo-ethyl, 4-methylbenzenesulfonyl, methylsulfonyl, 4-chlorobenzenesulfonyl, benzylsulfonyl, 4-methoxybenzenesulfonyl, benzenesulfonyl, 2,5-dimethoxybenzenesulfonyl, 2-cyanobenzenesulfonyl, thiophen-2-ylsulfonyl, 2-fluorobenzenesulfonyl, 2-trifluoromethoxybenzenesulfonyl, dimethylaminosulfonyl, benzoyl, pyridin-3-ylcarbonyl, 2,4,6-trichlorobenzenecarbonyl, tert-butylaminocarbonyl, dimethylaminocarbonylmethyl, 2-(4-methyl-piperazin-1-yl)-2-oxo-ethyl, 2-(4-pyridin-4-ylpiperazin-1-yl)ethanoyl, 2-[4-(2-methoxyphenyl)piperazin-1-yl]ethanoyl or 2-[4-(1,1-diphenylmethyl)piperazin-1-yl]ethanoyl,
 and the salts of these compounds.

5. Compounds of formula 1





in which

R1 is hydrogen or 1-4C-alkyl,

R2 is hydrogen or 1-4C-alkyl,

R3 represents a phenyl derivative of formulae (a) or (b)



wherein

R4 is 1-4C-alkoxy or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R5 is 1-8C-alkoxy, 3-7C-cycloalkoxy, 3-7C-cycloalkylmethoxy, or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R6 is 1-4C-alkoxy, 3-5C-cycloalkoxy, 3-5C-cycloalkylmethoxy, or 1-4C-alkoxy which is completely or predominantly substituted by fluorine,

R7 is 1-4C-alkyl and

R8 is hydrogen or 1-4C-alkyl,

or wherein

R7 and R8 together and with inclusion of the two carbon atoms, to which they are bonded, form a spiro-linked 5-, 6- or 7-membered hydrocarbon ring, optionally interrupted by an oxygen or sulphur atom,


R9 is hydrogen, 1-4C-alkyl, $-S(O)_2-R_{10}$, $-S(O)_2-(CH_2)_n-R_{11}$, $-(CH_2)_m-S(O)_2-R_{12}$, $-C(O)R_{13}$, $-C(O)-(CH_2)_n-R_{14}$, $-(CH_2)_m-C(O)-R_{15}$, Aryl1 or (Aryl2)-1-4C-alkyl,

R10 is 1-4C-alkyl, 5-dimethylaminonaphthalin-1-yl, $-N(R_{16})R_{17}$, phenyl or phenyl substituted by R18 and/or R19,

R11 is $-N(R_{16})R_{17}$,

R12 is $-N(R_{16})R_{17}$,

- R13 is 1-4C-alkyl, hydroxycarbonyl-1-4C-alkyl, phenyl, pyridyl, 4-ethyl-piperazin-2,3-dion-1-yl or -N(R16)R17,
 R14 is -N(R16)R17,
 R15 is -N(R16)R17, phenyl or phenyl substituted by R18 and/or R19 and/or R20,
 R16 and R17 are independent from each other hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-methyl, phenyl or phenyl substituted by R18 and/or R19 and/or R20, or R16 and R17 together and with inclusion of the nitrogen atom to which they are bonded, form a 4-morpholinyl-, 1-pyrrolidinyl-, 1-piperidinyl-, 1-hexahydroazepino- or a 1-piperazinyl-ring of formula (c)





wherein

- R21 is 1-4C-alkyl, pyrid-4-yl, pyrid-4-ylmethyl, dimethylamino-1-4C-alkyl, dimethylaminocarbonylmethyl, N-methyl-piperidin-4-yl, 4-morpholino-ethyl or tetrahydrofuran-2-ylmethyl,
 R18 is halogen, nitro, cyano, carboxyl, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, amino, mono- or di-1-4C-alkylamino, aminocarbonyl, 1-4C-alkylcarbonylamino or mono- or di-1-4C-alkylaminocarbonyl,
 R19 is halogen, amino, nitro, 1-4C-alkyl or 1-4C-alkoxy,
 R20 is halogen,
 Aryl1 is pyrimidin-2-yl, thieno-[2,3-d]pyrimidin-4-yl, 1-methyl-1H-pyrazolo-[3,4-d]pyrimidin-4-yl, thiazolyl, imidazolyl, furanyl, pyridyl, phenyl or phenyl substituted by R18 and/or R19,
 Aryl2 is pyridyl, phenyl, phenyl substituted by R18 and/or R19, 2-oxo-2H-chromen-7-yl or 4-(1,2,3-thiadiazol-4-yl)phenyl,
 n is an integer from 1 to 4,
 m is an integer from 1 to 4,
 and the salts of these compounds.

6. Compounds of formula 1 according to claim 5, in which

- R1 is hydrogen,
 R2 is hydrogen or 1-4C-alkyl,
 R3 represents a phenyl derivative of formulae (a) or (b)



wherein

R4 is 1-2C-alkoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,

R5 is 1-4C-alkoxy,

R6 is 1-2C-alkoxy or 1-2C-alkoxy which is completely or predominantly substituted by fluorine,

R7 is methyl and

R8 is hydrogen,

or wherein

R7 and R8 together and with inclusion of the two carbon atoms, to which they are bonded, form a spiro-linked cyclopentane, cyclohexane, tetrahydrofurane or tetrahydropyran ring,

R9 is hydrogen, 1-4C-alkyl, $-S(O)_2-R_{10}$, $-C(O)R_{13}$, $-(CH_2)_m-C(O)-R_{15}$, Aryl1 or (Aryl2)-1-4C-alkyl,

R10 is 1-4C-alkyl, 5-dimethylaminonaphthalin-1-yl or $-N(R_{16})R_{17}$,

R13 is 1-4C-alkyl, phenyl, 4-ethyl-piperazin-2,3-dion-1-yl or $-N(R_{16})R_{17}$,

R15 is $-N(R_{16})R_{17}$, phenyl or phenyl substituted by R18 and/or R19 and/or R20,

R16 and R17 are independent from each other hydrogen, 1-7C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-methyl, phenyl or phenyl substituted by R18 and/or R19 and/or R20, or R16 and R17 together and with inclusion of the nitrogen atom to which they are bonded, form a 4-morpholinyl-, 1-pyrrolidinyl-, 1-piperidinyl-, 1-hexahydroazepino- or a 4-methyl-piperazin-1-yl-ring,

R18 is halogen, nitro, cyano, 1-4C-alkyl, trifluoromethyl, 1-4C-alkoxy or 1-4C-alkoxycarbonyl,

R19 is halogen, amino, nitro, 1-4C-alkyl or 1-4C-alkoxy,

R20 is halogen,

Aryl1 is pyrimidin-2-yl, thieno-[2,3-d]pyrimidin-4-yl, 1-methyl-1H-pyrazolo-[3,4-d]pyrimidin-4-yl, pyridyl, phenyl or phenyl substituted by R18 and/or R19,

Aryl2 is pyridyl, phenyl, phenyl substituted by R18 and/or R19, 2-oxo-2H-chromen-7-yl or 4-(1,2,3-thiadiazol-4-yl)phenyl,

m is an integer from 1 to 2,



and the salts of these compounds.

7. Compounds of formula 1 according to claim 5, in which

R1 is hydrogen,

R2 is hydrogen or methyl,

R3 represents a phenyl derivative of formulae (a) or (b)




wherein

- R4 is 1-2C-alkoxy,
 R5 is 1-4C-alkoxy,
 R6 is 1-2C-alkoxy,
 R7 is methyl and
 R8 is hydrogen,
 R9 is hydrogen, $-S(O)_2-R_{10}$, $-C(O)R_{13}$, $-(CH_2)_m-C(O)-R_{15}$ or (Aryl2)-1-2C-alkyl,
 R10 is 1-4C-alkyl, 5-dimethylaminonaphthalin-1-yl or $-N(R_{16})R_{17}$,
 R13 is 1-4C-alkyl, 4-ethyl-piperazin-2,3-dion-1-yl or $-N(R_{16})R_{17}$,
 R15 is $-N(R_{16})R_{17}$,
 R16 and R17 are independent from each other hydrogen or 1-4C-alkyl, or R16 and R17 together and with inclusion of the nitrogen atom to which they are bonded, form a 4-morpholinyl ring, a 1-piperidinyl ring or a 4-methyl-piperazin-1-yl ring,
 Aryl2 is pyridyl or phenyl,
 m is 1,
 and the salts of these compounds.

8. Compounds of formula 1 according to claim 5, in which

- R1 is hydrogen,
 R2 is methyl,
 R3 represents a phenyl derivative of formula (a)



wherein

- R4 is methoxy,
 R5 is methoxy and
 R9 is acetyl, morpholin-4-ylcarbonyl, pyridin-3-ylmethyl, 4-ethyl-piperazin-2,3-dion-1-yl, 4-methylpiperazin-1-yl, 5-dimethylamino-naphthalene-1-sulfonyl or morpholin-4-yl-2-oxo-ethyl,
 and the salts of these compounds.

9. Compounds of formula 1 according to claim 1 or 5 for the treatment of diseases.

10. Pharmaceutical compositions containing one or more compounds of formula 1 according to claim 1 or 5 together with the usual pharmaceutical auxiliaries and/or carrier materials.

11. Use of compounds of formula 1 according to claim 1 or 5 for the preparation of pharmaceutical compositions for the treatment of airway disorders.
12. A method for treating an illness treatable by the administration of a PDE4 inhibitor in a patient comprising administering to said patient in need thereof a therapeutically effective amount of a compound of formula 1 as claimed in claim 1 or 5.
13. A method for treating airway disorders in a patient comprising administering to said patient a therapeutically effective amount of a compound of formula 1 as claimed in claim 1 or 5.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/08677

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/04 C07D401/14 A61K31/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 01 94319 A (BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH) 13 December 2001 (2001-12-13) page 64 -page 75; claims 1-10 ---	1-13
Y	WO 99 31090 A (BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH) 24 June 1999 (1999-06-24) page 34 -page 38; claims 1-10 ---	1-13
Y	EP 0 763 534 A (MERCK PATENT GMBH) 19 March 1997 (1997-03-19) cited in the application page 31 -page 34; claims 1-10 ---	1-13
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

29 October 2003

Date of mailing of the international search report

05/11/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kyriakakou, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/08677

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 92 06963 A (BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH) 30 April 1992 (1992-04-30) page 8 -page 10; claims 1-8 ---	1-13
A	EDDY SOTELO ET AL.: "Pyridazines. Part 25: Efficient and Selective....." TETRAHEDRON LETTERS., vol. 42, 2001, pages 8633-8636, XP004321511 ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM., NL ISSN: 0040-4039 the whole document ---	1-13
Y	WILLIAM J. COATES ET AL.: "1,4-Bis(3-Oxo-2,3-dihydropyridazin-6-yl)b enzene Analogues: Potent phosphodiesterase Inhibitors and Inodilators" JOURNAL OF MEDICINAL CHEMISTRY., vol. 33, no. 6, 1990, pages 1735-1741, XP002119282 AMERICAN CHEMICAL SOCIETY. WASHINGTON., US ISSN: 0022-2623 the whole document ---	1-13
Y	MARGARETHA VAN DER MEY ET AL.: "Novel Selective PDE4 Inhibitors. 1. Synthesis, Structure-Activity Relationships, and Molecular Modelling of 4-(3,4-Dimethoxyphenyl)-2H-phthalazin-1-on es and Analogues" JOURNAL OF MEDICINAL CHEMISTRY., vol. 44, no. 16, 2001, pages 2511-2522, XP002222950 AMERICAN CHEMICAL SOCIETY. WASHINGTON., US ISSN: 0022-2623 the whole document ---	1-13
Y	MARGARETHA VAN DER MEY ET AL.: "Novel Selective PDE4 Inhibitors.2. Synthesis and Structure-Activity Relationships of 4-Aryl-Substituted cis Tetra- and cis Hexahydrophthalazinones" JOURNAL OF MEDICINAL CHEMISTRY., vol. 44, no. 16, 2001, pages 2523-2535, XP002222951 AMERICAN CHEMICAL SOCIETY. WASHINGTON., US ISSN: 0022-2623 the whole document -----	1-13

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 03/08677

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 12 and 13 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/08677

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0194319	A	13-12-2001	AU 6233201 A	17-12-2001
			BR 0111440 A	03-06-2003
			CA 2411351 A1	13-12-2001
			CZ 20023997 A3	14-05-2003
			WO 0194319 A1	13-12-2001
			EP 1296956 A1	02-04-2003
			NO 20025811 A	04-02-2003
WO 9931090	A	24-06-1999	AU 753576 B2	24-10-2002
			AU 2270199 A	05-07-1999
			CA 2314111 A1	24-06-1999
			EE 200000335 A	15-10-2001
			WO 9931090 A1	24-06-1999
			EP 1042319 A1	11-10-2000
			HU 0004561 A2	28-03-2002
			JP 2002508368 T	19-03-2002
			PL 341239 A1	26-03-2001
			US 6380196 B1	30-04-2002
EP 763534	A	19-03-1997	DE 19533975 A1	20-03-1997
			AT 233258 T	15-03-2003
			AU 716113 B2	17-02-2000
			AU 6551796 A	20-03-1997
			BR 9603736 A	26-05-1998
			CA 2185397 A1	15-03-1997
			CN 1157287 A ,B	20-08-1997
			CZ 9602630 A3	18-03-1998
			DE 59610164 D1	03-04-2003
			DK 763534 T3	10-06-2003
			EP 0763534 A1	19-03-1997
			HU 9602511 A2	28-03-1997
			JP 9124611 A	13-05-1997
			NO 963852 A	17-03-1997
			PL 316070 A1	17-03-1997
			PT 763534 T	31-07-2003
			RU 2167159 C2	20-05-2001
			SK 110096 A3	06-08-1997
			US 5859008 A	12-01-1999
			ZA 9607766 A	26-03-1997
WO 9206963	A	30-04-1992	AT 168101 T	15-07-1998
			AU 8722991 A	20-05-1992
			CA 2094127 A1	17-04-1992
			DE 59109027 D1	13-08-1998
			WO 9206963 A1	30-04-1992
			EP 0553174 A1	04-08-1993
			JP 6501941 T	03-03-1994
			US 5376656 A	27-12-1994

CORRECTED VERSION

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
4 March 2004 (04.03.2004)

PCT

(10) International Publication Number
WO 2004/018451 A1

(51) International Patent Classification⁷: **C07D 401/04**,
401/14, A61K 31/50

(21) International Application Number:
PCT/EP2003/008677

(22) International Filing Date: 6 August 2003 (06.08.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
02017976.8 10 August 2002 (10.08.2002) EP

(71) Applicant (for all designated States except US): **ALTANA
PHARMA AG** [DE/DE]; Byk-Gulden-Str. 2, 78467 Kon-
stanz (DE).

(72) Inventors (for all designated States except CA, PH, US):
HATZELMANN, Armin; Alter Wall 3, 78467 Konstanz
(DE). **BARSIG, Johannes**; Bleichenweg 11, 78467 Kon-
stanz (DE). **MARX, Degenhard**; Obere Rente 15, 78345
Moos (DE). **KLEY, Hans-Peter**; Im Weinberg 3b, 78476
Allensbach (DE). **CHRISTIAANS, Johannes, A. M.**;
Zevenwouden 233, NL-3524 CR Utrecht (NL). **MENGE,**
Wiro, M., P., B.; Pontanuslaan 11, NL-6821 HM Arnhem
(NL).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **STERK, Geert, Jau**
[NL/NL]; Stadhouderslaan 38, NL-3583 JJ Utrecht (NL).

(74) Agent: **WILD, Robert**; Altana Pharma AG, Byk-Gulden-
Strasse 2, 78467 Konstanz (DE).

(81) Designated States (national): AE, AL, AU, BA, BR, CA,
CN, CO, DZ, EC, GE, HR, ID, IL, IN, IS, JP, KR, LT, LV,
MA, MK, MX, NO, NZ, PH, PL, SG, TN, UA, US, VN,
YU, ZA, ZW.

(84) Designated States (regional): Eurasian patent (AM, AZ,
BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE,
BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU,
IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AL, AU, BA, BR, CA, CN, CO, DZ, EC, GE, HR, ID, IL, IN,
IS, JP, KR, LT, LV, MA, MK, MX, NO, NZ, PH, PL, SG, TN,
UA, VN, YU, ZA, ZW, Eurasian patent (AM, AZ, BY, KG,
KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH,
CY, CZ, DE, DK, EE, ES, FI, FR, GR, GR, HU, IE, IT, LU,
MC, NL, PT, RO, SE, SI, SK, TR)
- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only

Published:

- with international search report

(48) Date of publication of this corrected version:

6 May 2004

(15) Information about Correction:

see PCT Gazette No. 19/2004 of 6 May 2004, Section II

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: PYRIDAZINONE-DERIVATIVES AS PDE4 INHIBITORS

(57) Abstract: The compounds of a generic formula (I), in which the substituents R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³⁰, R³¹, R³², R³³, R³⁴, R³⁵, R³⁶, R³⁷, R³⁸, R³⁹, R⁴⁰, R⁴¹, R⁴², R⁴³, R⁴⁴, R⁴⁵, R⁴⁶, R⁴⁷, R⁴⁸, R⁴⁹, R⁵⁰, R⁵¹, R⁵², R⁵³, R⁵⁴, R⁵⁵, R⁵⁶, R⁵⁷, R⁵⁸, R⁵⁹, R⁶⁰, R⁶¹, R⁶², R⁶³, R⁶⁴, R⁶⁵, R⁶⁶, R⁶⁷, R⁶⁸, R⁶⁹, R⁷⁰, R⁷¹, R⁷², R⁷³, R⁷⁴, R⁷⁵, R⁷⁶, R⁷⁷, R⁷⁸, R⁷⁹, R⁸⁰, R⁸¹, R⁸², R⁸³, R⁸⁴, R⁸⁵, R⁸⁶, R⁸⁷, R⁸⁸, R⁸⁹, R⁹⁰, R⁹¹, R⁹², R⁹³, R⁹⁴, R⁹⁵, R⁹⁶, R⁹⁷, R⁹⁸, R⁹⁹, R¹⁰⁰, R¹⁰¹, R¹⁰², R¹⁰³, R¹⁰⁴, R¹⁰⁵, R¹⁰⁶, R¹⁰⁷, R¹⁰⁸, R¹⁰⁹, R¹¹⁰, R¹¹¹, R¹¹², R¹¹³, R¹¹⁴, R¹¹⁵, R¹¹⁶, R¹¹⁷, R¹¹⁸, R¹¹⁹, R¹²⁰, R¹²¹, R¹²², R¹²³, R¹²⁴, R¹²⁵, R¹²⁶, R¹²⁷, R¹²⁸, R¹²⁹, R¹³⁰, R¹³¹, R¹³², R¹³³, R¹³⁴, R¹³⁵, R¹³⁶, R¹³⁷, R¹³⁸, R¹³⁹, R¹⁴⁰, R¹⁴¹, R¹⁴², R¹⁴³, R¹⁴⁴, R¹⁴⁵, R¹⁴⁶, R¹⁴⁷, R¹⁴⁸, R¹⁴⁹, R¹⁵⁰, R¹⁵¹, R¹⁵², R¹⁵³, R¹⁵⁴, R¹⁵⁵, R¹⁵⁶, R¹⁵⁷, R¹⁵⁸, R¹⁵⁹, R¹⁶⁰, R¹⁶¹, R¹⁶², R¹⁶³, R¹⁶⁴, R¹⁶⁵, R¹⁶⁶, R¹⁶⁷, R¹⁶⁸, R¹⁶⁹, R¹⁷⁰, R¹⁷¹, R¹⁷², R¹⁷³, R¹⁷⁴, R¹⁷⁵, R¹⁷⁶, R¹⁷⁷, R¹⁷⁸, R¹⁷⁹, R¹⁸⁰, R¹⁸¹, R¹⁸², R¹⁸³, R¹⁸⁴, R¹⁸⁵, R¹⁸⁶, R¹⁸⁷, R¹⁸⁸, R¹⁸⁹, R¹⁹⁰, R¹⁹¹, R¹⁹², R¹⁹³, R¹⁹⁴, R¹⁹⁵, R¹⁹⁶, R¹⁹⁷, R¹⁹⁸, R¹⁹⁹, R²⁰⁰, R²⁰¹, R²⁰², R²⁰³, R²⁰⁴, R²⁰⁵, R²⁰⁶, R²⁰⁷, R²⁰⁸, R²⁰⁹, R²¹⁰, R²¹¹, R²¹², R²¹³, R²¹⁴, R²¹⁵, R²¹⁶, R²¹⁷, R²¹⁸, R²¹⁹, R²²⁰, R²²¹, R²²², R²²³, R²²⁴, R²²⁵, R²²⁶, R²²⁷, R²²⁸, R²²⁹, R²³⁰, R²³¹, R²³², R²³³, R²³⁴, R²³⁵, R²³⁶, R²³⁷, R²³⁸, R²³⁹, R²⁴⁰, R²⁴¹, R²⁴², R²⁴³, R²⁴⁴, R²⁴⁵, R²⁴⁶, R²⁴⁷, R²⁴⁸, R²⁴⁹, R²⁵⁰, R²⁵¹, R²⁵², R²⁵³, R²⁵⁴, R²⁵⁵, R²⁵⁶, R²⁵⁷, R²⁵⁸, R²⁵⁹, R²⁶⁰, R²⁶¹, R²⁶², R²⁶³, R²⁶⁴, R²⁶⁵, R²⁶⁶, R²⁶⁷, R²⁶⁸, R²⁶⁹, R²⁷⁰, R²⁷¹, R²⁷², R²⁷³, R²⁷⁴, R²⁷⁵, R²⁷⁶, R²⁷⁷, R²⁷⁸, R²⁷⁹, R²⁸⁰, R²⁸¹, R²⁸², R²⁸³, R²⁸⁴, R²⁸⁵, R²⁸⁶, R²⁸⁷, R²⁸⁸, R²⁸⁹, R²⁹⁰, R²⁹¹, R²⁹², R²⁹³, R²⁹⁴, R²⁹⁵, R²⁹⁶, R²⁹⁷, R²⁹⁸, R²⁹⁹, R³⁰⁰, R³⁰¹, R³⁰², R³⁰³, R³⁰⁴, R³⁰⁵, R³⁰⁶, R³⁰⁷, R³⁰⁸, R³⁰⁹, R³¹⁰, R³¹¹, R³¹², R³¹³, R³¹⁴, R³¹⁵, R³¹⁶, R³¹⁷, R³¹⁸, R³¹⁹, R³²⁰, R³²¹, R³²², R³²³, R³²⁴, R³²⁵, R³²⁶, R³²⁷, R³²⁸, R³²⁹, R³³⁰, R³³¹, R³³², R³³³, R³³⁴, R³³⁵, R³³⁶, R³³⁷, R³³⁸, R³³⁹, R³⁴⁰, R³⁴¹, R³⁴², R³⁴³, R³⁴⁴, R³⁴⁵, R³⁴⁶, R³⁴⁷, R³⁴⁸, R³⁴⁹, R³⁵⁰, R³⁵¹, R³⁵², R³⁵³, R³⁵⁴, R³⁵⁵, R³⁵⁶, R³⁵⁷, R³⁵⁸, R³⁵⁹, R³⁶⁰, R³⁶¹, R³⁶², R³⁶³, R³⁶⁴, R³⁶⁵, R³⁶⁶, R³⁶⁷, R³⁶⁸, R³⁶⁹, R³⁷⁰, R³⁷¹, R³⁷², R³⁷³, R³⁷⁴, R³⁷⁵, R³⁷⁶, R³⁷⁷, R³⁷⁸, R³⁷⁹, R³⁸⁰, R³⁸¹, R³⁸², R³⁸³, R³⁸⁴, R³⁸⁵, R³⁸⁶, R³⁸⁷, R³⁸⁸, R³⁸⁹, R³⁹⁰, R³⁹¹, R³⁹², R³⁹³, R³⁹⁴, R³⁹⁵, R³⁹⁶, R³⁹⁷, R³⁹⁸, R³⁹⁹, R⁴⁰⁰, R⁴⁰¹, R⁴⁰², R⁴⁰³, R⁴⁰⁴, R⁴⁰⁵, R⁴⁰⁶, R⁴⁰⁷, R⁴⁰⁸, R⁴⁰⁹, R⁴¹⁰, R⁴¹¹, R⁴¹², R⁴¹³, R⁴¹⁴, R⁴¹⁵, R⁴¹⁶, R⁴¹⁷, R⁴¹⁸, R⁴¹⁹, R⁴²⁰, R⁴²¹, R⁴²², R⁴²³, R⁴²⁴, R⁴²⁵, R⁴²⁶, R⁴²⁷, R⁴²⁸, R⁴²⁹, R⁴³⁰, R⁴³¹, R⁴³², R⁴³³, R⁴³⁴, R⁴³⁵, R⁴³⁶, R⁴³⁷, R⁴³⁸, R⁴³⁹, R⁴⁴⁰, R⁴⁴¹, R⁴⁴², R⁴⁴³, R⁴⁴⁴, R⁴⁴⁵, R⁴⁴⁶, R⁴⁴⁷, R⁴⁴⁸, R⁴⁴⁹, R⁴⁵⁰, R⁴⁵¹, R⁴⁵², R⁴⁵³, R⁴⁵⁴, R⁴⁵⁵, R⁴⁵⁶, R⁴⁵⁷, R⁴⁵⁸, R⁴⁵⁹, R⁴⁶⁰, R⁴⁶¹, R⁴⁶², R⁴⁶³, R⁴⁶⁴, R⁴⁶⁵, R⁴⁶⁶, R⁴⁶⁷, R⁴⁶⁸, R⁴⁶⁹, R⁴⁷⁰, R⁴⁷¹, R⁴⁷², R⁴⁷³, R⁴⁷⁴, R⁴⁷⁵, R⁴⁷⁶, R⁴⁷⁷, R⁴⁷⁸, R⁴⁷⁹, R⁴⁸⁰, R⁴⁸¹, R⁴⁸², R⁴⁸³, R⁴⁸⁴, R⁴⁸⁵, R⁴⁸⁶, R⁴⁸⁷, R⁴⁸⁸, R⁴⁸⁹, R⁴⁹⁰, R⁴⁹¹, R⁴⁹², R⁴⁹³, R⁴⁹⁴, R⁴⁹⁵, R⁴⁹⁶, R⁴⁹⁷, R⁴⁹⁸, R⁴⁹⁹, R⁵⁰⁰, R⁵⁰¹, R⁵⁰², R⁵⁰³, R⁵⁰⁴, R⁵⁰⁵, R⁵⁰⁶, R⁵⁰⁷, R⁵⁰⁸, R⁵⁰⁹, R⁵¹⁰, R⁵¹¹, R⁵¹², R⁵¹³, R⁵¹⁴, R⁵¹⁵, R⁵¹⁶, R⁵¹⁷, R⁵¹⁸, R⁵¹⁹, R⁵²⁰, R⁵²¹, R⁵²², R⁵²³, R⁵²⁴, R⁵²⁵, R⁵²⁶, R⁵²⁷, R⁵²⁸, R⁵²⁹, R⁵³⁰, R⁵³¹, R⁵³², R⁵³³, R⁵³⁴, R⁵³⁵, R⁵³⁶, R⁵³⁷, R⁵³⁸, R⁵³⁹, R⁵⁴⁰, R⁵⁴¹, R⁵⁴², R⁵⁴³, R⁵⁴⁴, R⁵⁴⁵, R⁵⁴⁶, R⁵⁴⁷, R⁵⁴⁸, R⁵⁴⁹, R⁵⁵⁰, R⁵⁵¹, R⁵⁵², R⁵⁵³, R⁵⁵⁴, R⁵⁵⁵, R⁵⁵⁶, R⁵⁵⁷, R⁵⁵⁸, R⁵⁵⁹, R⁵⁶⁰, R⁵⁶¹, R⁵⁶², R⁵⁶³, R⁵⁶⁴, R⁵⁶⁵, R⁵⁶⁶, R⁵⁶⁷, R⁵⁶⁸, R⁵⁶⁹, R⁵⁷⁰, R⁵⁷¹, R⁵⁷², R⁵⁷³, R⁵⁷⁴, R⁵⁷⁵, R⁵⁷⁶, R⁵⁷⁷, R⁵⁷⁸, R⁵⁷⁹, R⁵⁸⁰, R⁵⁸¹, R⁵⁸², R⁵⁸³, R⁵⁸⁴, R⁵⁸⁵, R⁵⁸⁶, R⁵⁸⁷, R⁵⁸⁸, R⁵⁸⁹, R⁵⁹⁰, R⁵⁹¹, R⁵⁹², R⁵⁹³, R⁵⁹⁴, R⁵⁹⁵, R⁵⁹⁶, R⁵⁹⁷, R⁵⁹⁸, R⁵⁹⁹, R⁶⁰⁰, R⁶⁰¹, R⁶⁰², R⁶⁰³, R⁶⁰⁴, R⁶⁰⁵, R⁶⁰⁶, R⁶⁰⁷, R⁶⁰⁸, R⁶⁰⁹, R⁶¹⁰, R⁶¹¹, R⁶¹², R⁶¹³, R⁶¹⁴, R⁶¹⁵, R⁶¹⁶, R⁶¹⁷, R⁶¹⁸, R⁶¹⁹, R⁶²⁰, R⁶²¹, R⁶²², R⁶²³, R⁶²⁴, R⁶²⁵, R⁶²⁶, R⁶²⁷, R⁶²⁸, R⁶²⁹, R⁶³⁰, R⁶³¹, R⁶³², R⁶³³, R⁶³⁴, R⁶³⁵, R⁶³⁶, R⁶³⁷, R⁶³⁸, R⁶³⁹, R⁶⁴⁰, R⁶⁴¹, R⁶⁴², R⁶⁴³, R⁶⁴⁴, R⁶⁴⁵, R⁶⁴⁶, R⁶⁴⁷, R⁶⁴⁸, R⁶⁴⁹, R⁶⁵⁰, R⁶⁵¹, R⁶⁵², R⁶⁵³, R⁶⁵⁴, R⁶⁵⁵, R⁶⁵⁶, R⁶⁵⁷, R⁶⁵⁸, R⁶⁵⁹, R⁶⁶⁰, R⁶⁶¹, R⁶⁶², R⁶⁶³, R⁶⁶⁴, R⁶⁶⁵, R⁶⁶⁶, R⁶⁶⁷, R⁶⁶⁸, R⁶⁶⁹, R⁶⁷⁰, R⁶⁷¹, R⁶⁷², R⁶⁷³, R⁶⁷⁴, R⁶⁷⁵, R⁶⁷⁶, R⁶⁷⁷, R⁶⁷⁸, R⁶⁷⁹, R⁶⁸⁰, R⁶⁸¹, R⁶⁸², R⁶⁸³, R⁶⁸⁴, R⁶⁸⁵, R⁶⁸⁶, R⁶⁸⁷, R⁶⁸⁸, R⁶⁸⁹, R⁶⁹⁰, R⁶⁹¹, R⁶⁹², R⁶⁹³, R⁶⁹⁴, R⁶⁹⁵, R⁶⁹⁶, R⁶⁹⁷, R⁶⁹⁸, R⁶⁹⁹, R⁷⁰⁰, R⁷⁰¹, R⁷⁰², R⁷⁰³, R⁷⁰⁴, R⁷⁰⁵, R⁷⁰⁶, R⁷⁰⁷, R⁷⁰⁸, R⁷⁰⁹, R⁷¹⁰, R⁷¹¹, R⁷¹², R⁷¹³, R⁷¹⁴, R⁷¹⁵, R⁷¹⁶, R⁷¹⁷, R⁷¹⁸, R⁷¹⁹, R⁷²⁰, R⁷²¹, R⁷²², R⁷²³, R⁷²⁴, R⁷²⁵, R⁷²⁶, R⁷²⁷, R⁷²⁸, R⁷²⁹, R⁷³⁰, R⁷³¹, R⁷³², R⁷³³, R⁷³⁴, R⁷³⁵, R⁷³⁶, R⁷³⁷, R⁷³⁸, R⁷³⁹, R⁷⁴⁰, R⁷⁴¹, R⁷⁴², R⁷⁴³, R⁷⁴⁴, R⁷⁴⁵, R⁷⁴⁶, R⁷⁴⁷, R⁷⁴⁸, R⁷⁴⁹, R⁷⁵⁰, R⁷⁵¹, R⁷⁵², R⁷⁵³, R⁷⁵⁴, R⁷⁵⁵, R⁷⁵⁶, R⁷⁵⁷, R⁷⁵⁸, R⁷⁵⁹, R⁷⁶⁰, R⁷⁶¹, R⁷⁶², R⁷⁶³, R⁷⁶⁴, R⁷⁶⁵, R⁷⁶⁶, R⁷⁶⁷, R⁷⁶⁸, R⁷⁶⁹, R⁷⁷⁰, R⁷⁷¹, R⁷⁷², R⁷⁷³, R⁷⁷⁴, R⁷⁷⁵, R⁷⁷⁶, R⁷⁷⁷, R⁷⁷⁸, R⁷⁷⁹, R⁷⁸⁰, R⁷⁸¹, R⁷⁸², R⁷⁸³, R⁷⁸⁴, R⁷⁸⁵, R⁷⁸⁶, R⁷⁸⁷, R⁷⁸⁸, R⁷⁸⁹, R⁷⁹⁰, R⁷⁹¹, R⁷⁹², R⁷⁹³, R⁷⁹⁴, R⁷⁹⁵, R⁷⁹⁶, R⁷⁹⁷, R⁷⁹⁸, R⁷⁹⁹, R⁸⁰⁰, R⁸⁰¹, R⁸⁰², R⁸⁰³, R⁸⁰⁴, R⁸⁰⁵, R⁸⁰⁶, R⁸⁰⁷, R⁸⁰⁸, R⁸⁰⁹, R⁸¹⁰, R⁸¹¹, R⁸¹², R⁸¹³, R⁸¹⁴, R⁸¹⁵, R⁸¹⁶, R⁸¹⁷, R⁸¹⁸, R⁸¹⁹, R⁸²⁰, R⁸²¹, R⁸²², R⁸²³, R⁸²⁴, R⁸²⁵, R⁸²⁶, R⁸²⁷, R⁸²⁸, R⁸²⁹, R⁸³⁰, R⁸³¹, R⁸³², R⁸³³, R⁸³⁴, R⁸³⁵, R⁸³⁶, R⁸³⁷, R⁸³⁸, R⁸³⁹, R⁸⁴⁰, R⁸⁴¹, R⁸⁴², R⁸⁴³, R⁸⁴⁴, R⁸⁴⁵, R⁸⁴⁶, R⁸⁴⁷, R⁸⁴⁸, R⁸⁴⁹, R⁸⁵⁰, R⁸⁵¹, R⁸⁵², R⁸⁵³, R⁸⁵⁴, R⁸⁵⁵, R⁸⁵⁶, R⁸⁵⁷, R⁸⁵⁸, R⁸⁵⁹, R⁸⁶⁰, R⁸⁶¹, R⁸⁶², R⁸⁶³, R⁸⁶⁴, R⁸⁶⁵, R⁸⁶⁶, R⁸⁶⁷, R⁸⁶⁸, R⁸⁶⁹, R⁸⁷⁰, R⁸⁷¹, R⁸⁷², R⁸⁷³, R⁸⁷⁴, R⁸⁷⁵, R⁸⁷⁶, R⁸⁷⁷, R⁸⁷⁸, R⁸⁷⁹, R⁸⁸⁰, R⁸⁸¹, R⁸⁸², R⁸⁸³, R⁸⁸⁴, R⁸⁸⁵, R⁸⁸⁶, R⁸⁸⁷, R⁸⁸⁸, R⁸⁸⁹, R⁸⁹⁰, R⁸⁹¹, R⁸⁹², R⁸⁹³, R⁸⁹⁴, R⁸⁹⁵, R⁸⁹⁶, R⁸⁹⁷, R⁸⁹⁸, R⁸⁹⁹, R⁹⁰⁰, R⁹⁰¹, R⁹⁰², R⁹⁰³, R⁹⁰⁴, R⁹⁰⁵, R⁹⁰⁶, R⁹⁰⁷, R⁹⁰⁸, R⁹⁰⁹, R⁹¹⁰, R⁹¹¹, R⁹¹², R⁹¹³, R⁹¹⁴, R⁹¹⁵, R⁹¹⁶, R⁹¹⁷, R⁹¹⁸, R⁹¹⁹, R⁹²⁰, R⁹²¹, R⁹²², R⁹²³, R⁹²⁴, R⁹²⁵, R⁹²⁶, R⁹²⁷, R⁹²⁸, R⁹²⁹, R⁹³⁰, R⁹³¹, R⁹³², R⁹³³, R⁹³⁴, R⁹³⁵, R⁹³⁶, R⁹³⁷, R⁹³⁸, R⁹³⁹, R⁹⁴⁰, R⁹⁴¹, R⁹⁴², R⁹⁴³, R⁹⁴⁴, R⁹⁴⁵, R⁹⁴⁶, R⁹⁴⁷, R⁹⁴⁸, R⁹⁴⁹, R⁹⁵⁰, R⁹⁵¹, R⁹⁵², R⁹⁵³, R⁹⁵⁴, R⁹⁵⁵, R⁹⁵⁶, R⁹⁵⁷, R⁹⁵⁸, R⁹⁵⁹, R⁹⁶⁰, R⁹⁶¹, R⁹⁶², R⁹⁶³, R⁹⁶⁴, R⁹⁶⁵, R⁹⁶⁶, R⁹⁶⁷, R⁹⁶⁸, R⁹⁶⁹, R⁹⁷⁰, R⁹⁷¹, R⁹⁷², R⁹⁷³, R⁹⁷⁴, R⁹⁷⁵, R⁹⁷⁶, R⁹⁷⁷, R⁹⁷⁸, R⁹⁷⁹, R⁹⁸⁰, R⁹⁸¹, R⁹⁸², R⁹⁸³, R⁹⁸⁴, R⁹⁸⁵, R⁹⁸⁶, R⁹⁸⁷, R⁹⁸⁸, R⁹⁸⁹, R⁹⁹⁰, R⁹⁹¹, R⁹⁹², R⁹⁹³, R⁹⁹⁴, R⁹⁹⁵, R⁹⁹⁶, R⁹⁹⁷, R⁹⁹⁸, R⁹⁹⁹, R¹⁰⁰⁰, R¹⁰⁰¹, R¹⁰⁰², R¹⁰⁰³, R¹⁰⁰⁴, R¹⁰⁰⁵, R¹⁰⁰⁶, R¹⁰⁰⁷, R¹⁰⁰⁸, R¹⁰⁰⁹, R¹⁰¹⁰, R¹⁰¹¹, R¹⁰¹², R¹⁰¹³, R¹⁰¹⁴, R¹⁰¹⁵, R^{1016</}

0	For receiving Office use only	
0-1	International Application No.	
0-2	International Filing Date	
0-3	Name of receiving Office and "PCT International Application"	
0-4	Form - PCT/RO/101 PCT Request	
0-4-1	Prepared using	PCT-EASY Version 2.92 (updated 01.07.2003)
0-5	Petition The undersigned requests that the present international application be processed according to the Patent Cooperation Treaty	
0-6	Receiving Office (specified by the applicant)	European Patent Office (EPO) (RO/EP)
0-7	Applicant's or agent's file reference	1127WOORD01
I	Title of invention	NOVEL PYRIDAZINONE-DERIVATIVES
II	Applicant	
II-1	This person is:	applicant only
II-2	Applicant for	all designated States except US
II-4	Name	ALTANA PHARMA AG
II-5	Address:	Byk-Gulden-Str. 2 D-78467 Konstanz Germany
II-6	State of nationality	DE
II-7	State of residence	DE
II-8	Telephone No.	+49-(0) 7531-845226
II-9	Facsimile No.	+49-(0) 7531-845321
II-10	e-mail	DEKON.IPPA-DE@altanapharma.com
III-1	Applicant and/or Inventor	
III-1-1	This person is:	applicant and inventor
III-1-2	Applicant for	US only
III-1-4	Name (LAST, First)	STERK, Geert, Jan
III-1-5	Address:	Stadhouderslaan 38 NL-3583 JJ UTRECHT Netherlands
III-1-6	State of nationality	NL
III-1-7	State of residence	NL

III-2	Applicant and/or inventor	
III-2-1	This person is:	inventor only
III-2-3	Inventor for	EA: (AM AZ BY KG KZ MD RU TJ TM); EP: (AT BE BG CH&LI CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL PT RO SE SI SK TR); AE AL AU BA BR CN CO DZ EC GE HR ID IL IN IS JP KR LT LV MA MK MX NO NZ PH PL SG TN UA VN YU ZA ZW
III-2-4	Name (LAST, First)	HATZELMANN, Armin
III-2-5	Address:	Alter Wall 3 D-78467 Konstanz Germany
III-3	Applicant and/or inventor	
III-3-1	This person is:	inventor only
III-3-3	Inventor for	EA: (AM AZ BY KG KZ MD RU TJ TM); EP: (AT BE BG CH&LI CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL PT RO SE SI SK TR); AE AL AU BA BR CN CO DZ EC GE HR ID IL IN IS JP KR LT LV MA MK MX NO NZ PL SG TN UA VN YU ZA ZW
III-3-4	Name (LAST, First)	BARSIG, Johannes
III-3-5	Address:	Bleichenweg 11 D-78467 Konstanz Germany
III-4	Applicant and/or inventor	
III-4-1	This person is:	inventor only
III-4-3	Inventor for	EA: (AM AZ BY KG KZ MD RU TJ TM); EP: (AT BE BG CH&LI CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL PT RO SE SI SK TR); AE AL AU BA BR CN CO DZ EC GE HR ID IL IN IS JP KR LT LV MA MK MX NO NZ PH PL SG TN UA VN YU ZA ZW
III-4-4	Name (LAST, First)	MARX, Degenhard
III-4-5	Address:	Obere Reute 15 D-78345 Moos Germany

III-5	Applicant and/or inventor	
III-5-1	This person is:	inventor only
III-5-3	Inventor for	EA: (AM AZ BY KG KZ MD RU TJ TM); EP: (AT BE BG CH&LI CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL PT RO SE SI SK TR); AE AL AU BA BR CN CO DZ EC GE HR ID IL IN IS JP KR LT LV MA MK MX NO NZ PL SG TN UA VN YU ZA ZW
III-5-4	Name (LAST, First)	KLEY, Hans-Peter
III-5-5	Address:	Im Weinberg 3b D-78476 Allensbach Germany
III-6	Applicant and/or inventor	
III-6-1	This person is:	inventor only
III-6-3	Inventor for	EA: (AM AZ BY KG KZ MD RU TJ TM); EP: (AT BE BG CH&LI CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL PT RO SE SI SK TR); AE AL AU BA BR CN CO DZ EC GE HR ID IL IN IS JP KR LT LV MA MK MX NO NZ PL SG TN UA VN YU ZA ZW
III-6-4	Name (LAST, First)	CHRISTIAANS, Johannes, A. M.
III-6-5	Address:	Zevenwouden 233 NL-3524 CR UTRECHT Netherlands
III-7	Applicant and/or inventor	
III-7-1	This person is:	inventor only
III-7-3	Inventor for	EA: (AM AZ BY KG KZ MD RU TJ TM); EP: (AT BE BG CH&LI CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL PT RO SE SI SK TR); AE AL AU BA BR CN CO DZ EC GE HR ID IL IN IS JP KR LT LV MA MK MX NO NZ PH PL SG TN UA VN YU ZA ZW
III-7-4	Name (LAST, First)	MENGE, Wiro, M.P.B.
III-7-5	Address:	Pontanuslaan 11 NL-6821 HM ARNHEM Netherlands

IV-1	Agent or common representative; or address for correspondence The person identified below is hereby/has been appointed to act on behalf of the applicant(s) before the competent International Authorities as:	agent
IV-1-1	Name (LAST, First)	WILD, Robert
IV-1-2	Address:	c/o ALTANA Pharma AG Byk-Gulden-Straße 2 D-78467 Konstanz Germany
IV-1-3	Telephone No.	+49- (0) 7531-845226
IV-1-4	Facsimile No.	+49- (0) 7531-845321
IV-1-5	e-mail	DEKON.IPPA-DE@altanapharma.com
V	Designation of States	
V-1	Regional Patent (other kinds of protection or treatment, if any, are specified between parentheses after the designation(s) concerned)	EA: AM AZ BY KG KZ MD RU TJ TM and any other State which is a Contracting State of the Eurasian Patent Convention and of the PCT EP: AT BE BG CH&LI CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL PT RO SE SI SK TR and any other State which is a Contracting State of the European Patent Convention and of the PCT
V-2	National Patent (other kinds of protection or treatment, if any, are specified between parentheses after the designation(s) concerned)	AE AL AU BA BR CA CN CO DZ EC GE HR ID IL IN IS JP KR LT LV MA MK MX NO NZ PH PL SG TN UA US VN YU ZA ZW
V-5	Precautionary Designation Statement In addition to the designations made under items V-1, V-2 and V-3, the applicant also makes under Rule 4.9(b) all designations which would be permitted under the PCT except any designation(s) of the State(s) indicated under item V-6 below. The applicant declares that those additional designations are subject to confirmation and that any designation which is not confirmed before the expiration of 15 months from the priority date is to be regarded as withdrawn by the applicant at the expiration of that time limit.	
V-6	Exclusion(s) from precautionary designations	NONE
VI-1	Priority claim of earlier regional application	
VI-1-1	Filing date	10 August 2002 (10.08.2002)
VI-1-2	Number	02017976.8
VI-1-3	Regional Office	EP
VII-1	International Searching Authority Chosen	European Patent Office (EPO) (ISA/EP)

VII-2	Request to use results of earlier search; reference to that search		
VII-2-1	Date	28 November 2002 (28.11.2002)	
VII-2-2	Number	EP02017976	
VII-2-3	Country (or regional Office)	EP	
VIII	Declarations	Number of declarations	
VIII-1	Declaration as to the identity of the inventor	-	
VIII-2	Declaration as to the applicant's entitlement, as at the international filing date, to apply for and be granted a patent	1	
VIII-3	Declaration as to the applicant's entitlement, as at the international filing date, to claim the priority of the earlier application	1	
VIII-4	Declaration of inventorship (only for the purposes of the designation of the United States of America)	1	
VIII-5	Declaration as to non-prejudicial disclosures or exceptions to lack of novelty	-	



VIII-2-1	Declaration: Entitlement to apply for and be granted a patent Declaration as to the applicant's entitlement, as at the international filing date, to apply for and be granted a patent (Rules 4.17(ii) and 51bis.1(a)(ii)). In a case where the declaration under Rule 4.17(iv) is not appropriate: Name:	in relation to this international application ALTANA PHARMA AG is entitled to apply for and be granted a patent by virtue of the following:
VIII-2-1 (ii)		ALTANA PHARMA AG is entitled as employer of the inventor, HATZELMANN, Armin
VIII-2-1 (ii)		ALTANA PHARMA AG is entitled as employer of the inventor, BARSIG, Johannes
VIII-2-1 (ii)		ALTANA PHARMA AG is entitled as employer of the inventor, MARX, Degenhard
VIII-2-1 (ii)		ALTANA PHARMA AG is entitled as employer of the inventor, KLEY, Hans-Peter
VIII-2-1 (ii)		ALTANA PHARMA B. V. is entitled as employer of the inventor, CHRISTIAANS, Johannes A. M.
VIII-2-1 (ii)		ALTANA PHARMA B. V. is entitled as employer of the inventor, MENGE, Wiro
VIII-2-1 (ii)		ALTANA PHARMA B. V. is entitled as employer of the inventor, STERK, Geert, Jan
VIII-2-1 (iv)		an assignment from ALTANA PHARMA B. V. to ALTANA PHARMA AG, dated 22 July 2002 (22.07.2002)
VIII-2-1 (ix)	This declaration is made for the purposes of:	all designations except the designation of the United States of America

VIII-3-1	Declaration: Entitlement to claim priority Declaration as to the applicant's entitlement, as at the international filing date, to claim the priority of the earlier application specified below, where the applicant is not the applicant who filed the earlier application or where the applicant's name has changed since the filing of the earlier application (Rules 4.17(ii) and 51bis.1(a)(iii)): Name:	in relation to this international application ALTANA PHARMA AG is entitled to claim priority of earlier application No. EP 02017976.8 by virtue of the following:
VIII-3-1 (i)		the applicant is the inventor of the subject matter for which protection was sought by way of the earlier application
VIII-3-1 (iv)		an assignment from ALTANA PHARMA B. V. to ALTANA PHARMA AG, dated 22 July 2002 (22.07.2002)
VIII-3-1 (ix)	This declaration is made for the purposes of:	all designations

VIII-4-1	<p>Declaration: Inventorship (only for the purposes of the designation of the United States of America) Declaration of Inventorship (Rules 4.17(iv) and 51bis.1(a)(iv)) for the purposes of the designation of the United States of America:</p>	<p>I hereby declare that I believe I am the original, first and sole (if only one inventor is listed below) or joint (if more than one inventor is listed below) inventor of the subject matter which is claimed and for which a patent is sought.</p> <p>This declaration is directed to the international application of which it forms a part (if filing declaration with application).</p> <p>I hereby declare that my residence, mailing address, and citizenship are as stated next to my name.</p> <p>I hereby state that I have reviewed and understand the contents of the above-identified international application, including the claims of said application. I have identified in the request of said application, in compliance with PCT Rule 4.10, any claim to foreign priority, and I have identified below, under the heading "Prior Applications," by application number, country or Member of the World Trade Organization, day, month and year of filing, any application for a patent or inventor's certificate filed in a country other than the United States of America, including any PCT international application designating at least one country other than the United States of America, having a filing date before that of the application on which foreign priority is claimed.</p>
VIII-4-1 -1	Prior applications:	

		<p>I hereby acknowledge the duty to disclose information that is known by me to be material to patentability as defined by 37 C.F.R. § 1.56, including for continuation-in-part applications, material information which became available between the filing date of the prior application and the PCT international filing date of the continuation-in-part application.</p> <p>I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.</p>
VIII-4-1 -1-1	Name:	STERK, Geert, Jan
VIII-4-1 -1-2	Residence: (city and either US State, if applicable, or country)	JJ UTRECHT, Netherlands
VIII-4-1 -1-3	Mailing address:	Stadhouderslaan 38
VIII-4-1 -1-4	Citizenship:	NL
VIII-4-1 -1-5	Inventor's Signature: (If not contained in the request, or if declaration is corrected or added under Rule 26ter after the filing of the international application. The signature must be that of the inventor, not that of the agent)	16-08-2003 <i>G/Blak</i>
VIII-4-1 -1-6	Date: (of signature which is not contained in the request, or of the declaration that is corrected or added under Rule 26ter after the filing of the international application)	

Original (for SUBMISSION) - printed on 05.08.2003 10:07 AM

IX	Check list	number of sheets	electronic file(s) attached
IX-1	Request (including declaration sheets)	10	--
IX-2	Description	37	--
IX-3	Claims	10	--
IX-4	Abstract	1	EZABST00.TXT
IX-5	Drawings	0	--
IX-7	TOTAL	58	
	Accompanying items	paper document(s) attached	electronic file(s) attached
IX-8	Fee calculation sheet	✓	--
IX-13	Priority document(s)	Item(s) VI-1	--
IX-17	PCT-EASY diskette	--	Diskette
IX-19	Figure of the drawings which should accompany the abstract		
IX-20	Language of filing of the international application	English	
X-1	Signature of applicant, agent or common representative		
X-1-1	Name (LAST, First)	WILD, Robert	
X-2	Signature of applicant, agent or common representative		
X-2-1	Name (LAST, First)	STERK, Geert, Jan	

FOR RECEIVING OFFICE USE ONLY

10-1	Date of actual receipt of the purported international application	
10-2	Drawings:	
10-2-1	Received	
10-2-2	Not received	
10-3	Corrected date of actual receipt due to later but timely received papers or drawings completing the purported international application	
10-4	Date of timely receipt of the required corrections under PCT Article 11(2)	
10-5	International Searching Authority	ISA/EP
10-6	Transmittal of search copy delayed until search fee is paid	

FOR INTERNATIONAL BUREAU USE ONLY

11-1	Date of receipt of the record copy by the International Bureau	
------	--	--